



Part VI
Practice Test 2

AP[®] Chemistry Exam

SECTION I: Multiple-Choice Questions

DO NOT OPEN THIS BOOKLET UNTIL YOU ARE TOLD TO DO SO.**At a Glance****Total Time**

1 hour and 30 minutes

Number of Questions

60

Percent of Total Grade

50%

Writing Instrument

Pencil required

Instructions

Section I of this examination contains 60 multiple-choice questions. Fill in only the ovals for numbers 1 through 60 on your answer sheet.

CALCULATORS MAY NOT BE USED IN THIS PART OF THE EXAMINATION.

Indicate all of your answers to the multiple-choice questions on the answer sheet. No credit will be given for anything written in this exam booklet, but you may use the booklet for notes or scratch work. After you have decided which of the suggested answers is best, completely fill in the corresponding oval on the answer sheet. Give only one answer to each question. If you change an answer, be sure that the previous mark is erased completely. Here is a sample question and answer.

Sample Question

Chicago is a
(A) state
(B) city
(C) country
(D) continent

Sample Answer

(A) ● (C) (D)

Use your time effectively, working as quickly as you can without losing accuracy. Do not spend too much time on any one question. Go on to other questions and come back to the ones you have not answered if you have time. It is not expected that everyone will know the answers to all the multiple-choice questions.

About Guessing

Many candidates wonder whether or not to guess the answers to questions about which they are not certain. Multiple-choice scores are based on the number of questions answered correctly. Points are not deducted for incorrect answers, and no points are awarded for unanswered questions. Because points are not deducted for incorrect answers, you are encouraged to answer all multiple-choice questions. On any questions you do not know the answer to, you should eliminate as many choices as you can, and then select the best answer among the remaining choices.

GO ON TO THE NEXT PAGE.

ADVANCED PLACEMENT CHEMISTRY EQUATIONS AND CONSTANTS

Throughout the test the following symbols have the definitions specified unless otherwise noted.

L, mL = liter(s), milliliter(s)
 g = gram(s)
 nm = nanometer(s)
 atm = atmosphere(s)

mm Hg = millimeters of mercury
 J, kJ = joule(s), kilojoule(s)
 V = volt(s)
 mol = mole(s)

ATOMIC STRUCTURE

$$E = h\nu$$

$$c = \lambda\nu$$

E = energy

ν = frequency

λ = wavelength

Planck's constant, $h = 6.626 \times 10^{-34}$ J s

Speed of light, $c = 2.998 \times 10^8$ m s⁻¹

Avogadro's number = 6.022×10^{23} mol⁻¹

Electron charge, $e = -1.602 \times 10^{-19}$ coulomb

EQUILIBRIUM

$$K_c = \frac{[C]^c [D]^d}{[A]^a [B]^b}, \text{ where } a A + b B \rightleftharpoons c C + d D$$

$$K_p = \frac{(P_C)^c (P_D)^d}{(P_A)^a (P_B)^b}$$

$$K_a = \frac{[H^+][A^-]}{[HA]}$$

$$K_b = \frac{[OH^-][HB^+]}{[B]}$$

$$K_w = [H^+][OH^-] = 1.0 \times 10^{-14} \text{ at } 25^\circ\text{C}$$

$$= K_a \times K_b$$

$$\text{pH} = -\log[H^+], \text{ pOH} = -\log[OH^-]$$

$$14 = \text{pH} + \text{pOH}$$

$$\text{pH} = \text{p}K_a + \log \frac{[A^-]}{[HA]}$$

$$\text{p}K_a = -\log K_a, \text{ p}K_b = -\log K_b$$

Equilibrium Constants

K_c (molar concentrations)

K_p (gas pressures)

K_a (weak acid)

K_b (weak base)

K_w (water)

KINETICS

$$\ln[A]_t - \ln[A]_0 = -kt$$

$$\frac{1}{[A]_t} - \frac{1}{[A]_0} = kt$$

$$t_{1/2} = \frac{0.693}{k}$$

k = rate constant

t = time

$t_{1/2}$ = half-life

GO ON TO THE NEXT PAGE.

GASES, LIQUIDS, AND SOLUTIONS

$$PV = nRT$$

$$P_A = P_{\text{total}} \times X_A, \text{ where } X_A = \frac{\text{moles A}}{\text{total moles}}$$

$$P_{\text{total}} = P_A + P_B + P_C + \dots$$

$$n = \frac{m}{M}$$

$$K = ^\circ\text{C} + 273$$

$$D = \frac{m}{V}$$

$$KE \text{ per molecule} = \frac{1}{2}mv^2$$

Molarity, M = moles of solute per liter of solution

$$A = abc$$

P = pressure

V = volume

T = temperature

n = number of moles

m = mass

M = molar mass

D = density

KE = kinetic energy

v = velocity

A = absorbance

a = molar absorptivity

b = path length

c = concentration

$$\begin{aligned} \text{Gas constant, } R &= 8.314 \text{ J mol}^{-1} \text{ K}^{-1} \\ &= 0.08206 \text{ L atm mol}^{-1} \text{ K}^{-1} \\ &= 62.36 \text{ L torr mol}^{-1} \text{ K}^{-1} \end{aligned}$$

$$\begin{aligned} 1 \text{ atm} &= 760 \text{ mm Hg} \\ &= 760 \text{ torr} \end{aligned}$$

STP = 273.15 K and 1.0 atm

Ideal gas at STP = 22.4 L mol⁻¹

THERMOCHEMISTRY/ ELECTROCHEMISTRY

$$q = mc\Delta T$$

$$\Delta S^\circ = \sum S^\circ \text{ products} - \sum S^\circ \text{ reactants}$$

$$\Delta H^\circ = \sum \Delta H_f^\circ \text{ products} - \sum \Delta H_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \sum \Delta G_f^\circ \text{ products} - \sum \Delta G_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

$$= -RT \ln K$$

$$= -nFE^\circ$$

$$I = \frac{q}{t}$$

q = heat

m = mass

c = specific heat capacity

T = temperature

S° = standard entropy

H° = standard enthalpy

G° = standard free energy

n = number of moles

E° = standard reduction potential

I = current (amperes)

q = charge (coulombs)

t = time (seconds)

Faraday's constant, F = 96,485 coulombs per mole of electrons


$$1 \text{ volt} = \frac{1 \text{ joule}}{1 \text{ coulomb}}$$

GO ON TO THE NEXT PAGE.

1. A compound is made up of entirely silicon and oxygen atoms. If there are 14.0 g of silicon and 32.0 g of oxygen present, what is the empirical formula of the compound?


(A) SiO_2
 (B) SiO_4
 (C) Si_2O
 (D) Si_2O_3

2.



The volume of a gas is charted over time, giving the above results. Which of the following options provides a possible explanation of what was happening to the gas during each phase of the graph?

- (A) During phase I, the temperature decreased while the pressure increased. During phase II, the temperature was held constant as the pressure decreased.
- (B) During phase I, the temperature increased while the pressure was held constant. During phase II, the temperature and pressure both decreased.
- (C) During phase I, the temperature was held constant while the pressure increased. During phase II, the temperature and pressure both decreased.
- (D) During phase I, the temperature and pressure both increased. During phase II, the temperature was held constant while the pressure decreased.
3. A solution of sulfurous acid, H_2SO_3 , is present in an aqueous solution. Which of the following represents the concentrations of three different ions in solution?
- (A) $[\text{SO}_3^{2-}] > [\text{HSO}_3^-] > [\text{H}_2\text{SO}_3]$
 (B) $[\text{H}_2\text{SO}_3] > [\text{HSO}_3^-] > [\text{SO}_3^{2-}]$
 (C) $[\text{H}_2\text{SO}_3] > [\text{HSO}_3^-] = [\text{SO}_3^{2-}]$
 (D) $[\text{SO}_3^{2-}] = [\text{HSO}_3^-] > [\text{H}_2\text{SO}_3]$




The above experiment was performed several times, and the following data was gathered:

Trial	$[\text{NO}]_{\text{init}}$ (M)	$[\text{Br}_2]_{\text{init}}$ (M)	Initial Rate of Reaction (M/min)
1	0.20 M	0.10 M	5.20×10^{-3}
2	0.20 M	0.20 M	1.04×10^{-2}
3	0.40 M	0.10 M	2.08×10^{-2}

What is the rate law for this reaction?

(A) Rate = $k[\text{NO}][\text{Br}_2]^2$
 (B) Rate = $k[\text{NO}]^2[\text{Br}_2]^2$
 (C) Rate = $k[\text{NO}][\text{Br}_2]$
 (D) Rate = $k[\text{NO}]^2[\text{Br}_2]$



Which of the following statements accurately describes the above reaction?

(A) The entropy of the reactants exceeds that of the products.
 (B) $\text{H}_2\text{O}(l)$ will always be the limiting reagent.
 (C) This reaction is never thermodynamically favored.
 (D) The temperature of the surroundings will increase as this reaction progresses.


Use the following information to answer questions 6-9.

20.0 mL of 1.0 M Na_2CO_3 is placed in a beaker and titrated with a solution of 1.0 M $\text{Ca}(\text{NO}_3)_2$, resulting in the creation of a precipitate.


6. How much $\text{Ca}(\text{NO}_3)_2$ must be added to reach the equivalence point?
- (A) 10.0 mL
 (B) 20.0 mL
 (C) 30.0 mL
 (D) 40.0 mL

GO ON TO THE NEXT PAGE.


7. Which of the following diagrams correctly shows the species present in the solution in significant amounts at the equivalence point?




(A)



(B)



(C)



(D)

8. What will happen to the conductivity of the solution after additional $\text{Ca}(\text{NO}_3)_2$ is added past the equivalence point?
- (A) The conductivity will increase as additional ions are being added to the solution.
- (B) The conductivity will stay constant as the precipitation reaction has gone to completion.
- (C) The conductivity will decrease as the solution will be diluted with the addition of additional $\text{Ca}(\text{NO}_3)_2$.
- (D) The conductivity will stay constant as equilibrium has been established.
9. If the experiment were repeated and the Na_2CO_3 was diluted to 40.0 mL with distilled water prior to the titration, how would that affect the volume of $\text{Ca}(\text{NO}_3)_2$ needed to reach the equivalence point?
- (A) It would be cut in half.
- (B) It would decrease by a factor of 1.5.
- (C) It would double.
- (D) It would not change.




2.0 mol of $\text{CO}(g)$ and 2.0 mol of $\text{O}_2(g)$ are pumped into a rigid, evacuated 4.0-L container, where they react to form $\text{CO}_2(g)$. Which of the following values does NOT represent a potential set of concentrations for each gas at a given point during the reaction?

	CO	O ₂	CO ₂
(A)	0.5	0.5	0
(B)	0	0.25	0.5
(C)	0.25	0.25	0.5
(D)	0.25	0.38	0.25

11. Neutral atoms of chlorine are bombarded by high-energy photons, causing the ejection of electrons from the various filled subshells. Electrons originally from which subshell would have the highest velocity after being ejected?
- (A) $1s$
- (B) $2p$
- (C) $3p$
- (D) $3d$
12. A sample of oxygen gas at 50°C is heated, reaching a final temperature of 100°C . Which statement best describes the behavior of the gas molecules?
- (A) Their velocity increases by a factor of two.
- (B) Their velocity increases by a factor of four.
- (C) Their kinetic energy increases by a factor of 2.
- (D) Their kinetic energy increases by a factor of less than 2.
13. The average mass, in grams, of one mole of carbon atoms is equal to
- (A) the average mass of a single carbon atom, measured in amu
- (B) the ratio of the number of carbon atoms to the mass of a single carbon atom
- (C) the number of carbon atoms in one amu of carbon
- (D) the mass, in grams, of the most abundant isotope of carbon

GO ON TO THE NEXT PAGE.


14.



The diagram above best represents which type of reaction?

- (A) Acid/base
 (B) Oxidation/reduction
 (C) Precipitation
 (D) Decomposition
15. Which of the following is true for all bases?
- (A) All bases donate OH^- ions into solution.
 (B) Only strong bases create solutions in which OH^- ions are present.
 (C) Only strong bases are good conductors when dissolved in solution.
 (D) For weak bases, the concentration of the OH^- ions exceeds the concentration of the base in the solution.

Use the following information to answer questions 16-18.




In the above reaction, a piece of solid nickel is added to a solution of potassium dichromate.


16. Which species is being oxidized and which is being reduced?
- | | Oxidized | Reduced |
|-----|----------------------------------|----------------------------------|
| (A) | $\text{Cr}_2\text{O}_7^{2-}(aq)$ | $\text{Ni}(s)$ |
| (B) | $\text{Cr}^{3+}(aq)$ | $\text{Ni}^{2+}(aq)$ |
| (C) | $\text{Ni}(s)$ | $\text{Cr}_2\text{O}_7^{2-}(aq)$ |
| (D) | $\text{Ni}^{2+}(aq)$ | $\text{Cr}^{3+}(aq)$ |
17. How many moles of electrons are transferred when 1 mole of potassium dichromate is mixed with 3 mol of nickel?
- (A) 2 moles of electrons
 (B) 3 moles of electrons
 (C) 5 moles of electrons
 (D) 6 moles of electrons

18. How does the pH of the solution change as the reaction progresses?
- (A) It increases until the solution becomes basic.
 (B) It increases, but the solution remains acidic.
 (C) It decreases until the solution becomes basic.
 (D) It decreases, but the solution remains acidic.


19. A sample of an unknown chloride compound was dissolved in water, and then titrated with excess $\text{Pb}(\text{NO}_3)_2$ to create a precipitate. After drying, it is determined there are 0.0050 mol of precipitate present. What mass of chloride is present in the original sample?
- (A) 0.177 g
 (B) 0.355 g
 (C) 0.522 g
 (D) 0.710 g
20. A photoelectron spectra for which of the following atoms would show peaks at exactly three different binding energies?




(A)



(B)



(C)



(D)

21. The bond length between any two nonmetal atoms is achieved under which of the following conditions?
- (A) Where the energy of interaction between the atoms is at its minimum value
 (B) Where the nuclei of each atom exhibits the strongest attraction to the electrons of the other atom
 (C) The point at which the attractive and repulsive forces between the two atoms are equal
 (D) The closest point at which a valence electron from one atom can transfer to the other atom

GO ON TO THE NEXT PAGE.

22. Hydrogen fluoride, HF, is a liquid at 15°C. All other hydrogen halides (represented by HX, where X is any other halogen) are gases at the same temperature. Why?
- (A) Fluorine has a very high electronegativity; therefore, the H–F bond is stronger than any other H–X bond.
- (B) HF is smaller than any other H–X molecule; therefore, it exhibits stronger London dispersion forces.
- (C) The dipoles in a HF molecule exhibit a particularly strong attraction force to the dipoles in other HF molecules.
- (D) The H–F bond is the most ionic in character compared to all other hydrogen halides.

23.


	Initial pH	pH after NaOH addition
Acid 1	3.0	3.5
Acid 2	3.0	5.0

Two different acids with identical pH are placed in separate beakers. Identical portions of NaOH are added to each beaker, and the resulting pH is indicated in the table above. What can be determined about the strength of each acid?

- (A) Acid 1 is a strong acid and acid 2 is a weak acid because acid 1 resists change in pH more effectively.
- (B) Acid 1 is a strong acid and acid 2 is a weak acid because the NaOH is more effective at neutralizing acid 2.
- (C) Acid 1 is a weak acid and acid 2 is a strong acid because the concentration of the weak acid must be significantly greater to have the same pH as the strong acid.
- (D) Acid 1 is a weak acid and acid 2 is a strong acid because the concentration of the hydrogen ions will be greater in acid 2 after the NaOH addition.
24. A stock solution of 12.0 M sulfuric acid is made available. What is the best procedure to make up 100.0 mL of 4.0 M sulfuric acid using the stock solution and water prior to mixing?
- (A) Add 33.3 mL of water to the flask, and then add 66.7 mL of 12.0 M acid.
- (B) Add 33.3 mL of 12.0 M acid to the flask, and then dilute it with 66.7 mL of water.
- (C) Add 67.7 mL of 12.0 M acid to the flask, and then dilute it with 33.3 mL of water.
- (D) Add 67.7 mL of water to the flask, and then add 33.3 mL of 12.0 M acid.

Use the following data to answer questions 25-29.


The enthalpy values for several reactions are as follows:



25. In which of the reactions does the amount of energy released by the formation of bonds in the products exceed the amount of energy necessary to break the bonds of the reactants by the greatest amount?
- (A) Reaction I
 (B) Reaction II
 (C) Reaction III
 (D) Reaction IV
26. In which of the reactions is the value for ΔS the most positive?
- (A) Reaction I
 (B) Reaction II
 (C) Reaction III
 (D) Reaction IV
27. Regarding reaction I, how would the addition of a catalyst affect the enthalpy and entropy changes for this reaction?
- | | | |
|-----|-----------|-----------|
| | Enthalpy | Entropy |
| (A) | Decrease | Decrease |
| (B) | Decrease | No Change |
| (C) | No Change | Decrease |
| (D) | No Change | No Change |

GO ON TO THE NEXT PAGE.

28.



Regarding reaction II, to achieve the products present in the above diagram how many moles of each reactant must be present prior to the reaction?

- (A) 1.0 mol of CH_4 and 2.0 mol of H_2O
 (B) 2.0 mol of CH_4 and 2.0 mol of H_2O
 (C) 2.0 mol of CH_4 and 3.0 mol of H_2O
 (D) 3.0 mol of CH_4 and 2.0 mol of H_2O

29. Regarding reaction IV, how much heat is absorbed or released when 2.0 mol of $\text{CH}_4(g)$ reacts with 2.0 mol of $\text{O}_2(g)$?

- (A) 890 kJ of heat is released.
 (B) 890 kJ of heat is absorbed.
 (C) 1780 kJ of heat is released.
 (D) 1780 kJ of heat is absorbed.

30. London dispersion forces are caused by

- (A) temporary dipoles created by the position of electrons around the nuclei in a molecule
 (B) the three-dimensional intermolecular bonding present in all covalent substances
 (C) the uneven electron-to-proton ratio found on individual atoms of a molecule
 (D) the electronegativity differences between the different atoms in a molecule

31. What is the general relationship between temperature and entropy for diatomic gases?

- (A) They are completely independent of each other; temperature has no effect on entropy.
 (B) There is a direct relationship, because at higher temperatures there is an increase in energy dispersal.
 (C) There is an inverse relationship, because at higher temperatures substances are more likely to be in a gaseous state.
 (D) It depends on the specific gas and the strength of the intermolecular forces between individual molecules.

32. Mixing equimolar amounts of which of the following solutions would create a buffer with a pH between 4 and 5?

$$K_a \text{ for } \text{HC}_3\text{H}_2\text{O}_2 = 1.75 \times 10^{-5}$$


$$K_a \text{ for } \text{HPO}_4^{2-} = 4.8 \times 10^{-13}$$

- (A) H_2SO_4 and H_2PO_4^-
 (B) HPO_4^{2-} and Na_3PO_4
 (C) $\text{HC}_3\text{H}_2\text{O}_2$ and $\text{NaC}_3\text{H}_2\text{O}_2$
 (D) NaOH and $\text{HC}_2\text{H}_3\text{O}_2$

33. A solution contains a mixture of four different compounds: $\text{KCl}(aq)$, $\text{Fe}(\text{NO}_3)_3(aq)$, $\text{MgSO}_4(aq)$, and $\text{N}_2\text{H}_4(aq)$. Which of these compounds would be easiest to separate via distillation?

- (A) $\text{KCl}(aq)$
 (B) $\text{Fe}(\text{NO}_3)_3(aq)$
 (C) $\text{MgSO}_4(aq)$
 (D) $\text{N}_2\text{H}_4(aq)$

34.



Identify the three gases represented on the Maxwell-Boltzmann diagram above. Assume all gases are at the same temperature.

- | | I | II | III |
|-----|--------------|--------------|--------------|
| (A) | H_2 | N_2 | F_2 |
| (B) | H_2 | F_2 | N_2 |
| (C) | F_2 | N_2 | H_2 |
| (D) | N_2 | F_2 | H_2 |

GO ON TO THE NEXT PAGE.

Section I

35. A sample of solid MgCl_2 would be most soluble in which of the following solutions?
- (A) $\text{LiOH} (aq)$
 (B) $\text{CBr}_4 (aq)$
 (C) $\text{Mg}(\text{NO}_3)_2 (aq)$
 (D) $\text{AlCl}_3 (aq)$
36. Most transition metals share a common oxidation state of +2. Which of the following best explains why?
- (A) Transition metals all have a minimum of two unpaired electrons.
 (B) Transition metals have unstable configurations and are very reactive.
 (C) Transition metals tend to gain electrons when reacting with other elements.
 (D) Transition metals will lose their outermost s -block electrons when forming bonds.
37. $2\text{Ag}^+ (aq) + \text{Fe} (s) \rightarrow 2\text{Ag} (s) + \text{Fe}^{2+} (aq)$

Which of the following would cause an increase in potential in the voltaic cell described by the above reaction?


- (A) Increasing $[\text{Fe}^{2+}]$
 (B) Adding more $\text{Fe} (s)$
 (C) Decreasing $[\text{Fe}^{2+}]$
 (D) Removing some $\text{Fe} (s)$
- Use the following information to answer questions 38-41.

Consider the Lewis structures for the following molecules:



38. Which molecule would have the shortest bonds?
- (A) CO_2
 (B) CO_3^{2-}
 (C) NO_2^-
 (D) NO_3^-
39. Which molecules are best represented by multiple resonance structures?
- (A) CO_2 and CO_3^{2-}
 (B) NO_2^- and NO_3^-
 (C) CO_3^{2-} and NO_3^-
 (D) CO_3^{2-} , NO_2^- , and NO_3^-


40. Which molecule or molecules exhibit sp^2 hybridization around the central atom?
- (A) CO_2 and CO_3^{2-}
 (B) NO_2^- and NO_3^-
 (C) CO_3^{2-} and NO_3^-
 (D) CO_3^{2-} , NO_2^- , and NO_3^-
41. Which molecule would have the smallest bond angle between terminal atoms?
- (A) CO_2
 (B) CO_3^{2-}
 (C) NO_2^-
 (D) NO_3^-




Increasing the temperature of the above reaction will increase the rate of reaction. Which of the following is NOT a reason that increased temperature increases reaction rate?

- (A) The reactants will be more likely to overcome the activation energy.
 (B) The number of collisions between reactant molecules will increase.
 (C) A greater distribution of reactant molecules will have high velocities.
 (D) Alternate reaction pathways become available at higher temperatures.


43. Which of the following diagrams best represents what is happening on a molecular level when NaCl dissolves in water?




(A)



(B)



(C)



(D)

GO ON TO THE NEXT PAGE.

44. Nitrous acid, HNO_2 , has a $\text{p}K_a$ value of 3.3. If a solution of nitrous acid is found to have a pH of 4.2, what can be said about the concentration of the conjugate acid/base pair found in solution?


- (A) $[\text{HNO}_2] > [\text{NO}_2^-]$
 (B) $[\text{NO}_2^-] > [\text{HNO}_2]$
 (C) $[\text{H}_2\text{NO}_2^+] > [\text{HNO}_2]$
 (D) $[\text{HNO}_2] > [\text{H}_2\text{NO}_2^+]$

45. Which of the following processes is an irreversible reaction?

- (A) $\text{CH}_4(g) + \text{O}_2(g) \rightarrow \text{CO}_2(g) + \text{H}_2\text{O}(l)$
 (B) $\text{HCN}(aq) + \text{H}_2\text{O}(l) \rightarrow \text{CN}^-(aq) + \text{H}_3\text{O}^+(aq)$
 (C) $\text{Al}(\text{NO}_3)_3(s) \rightarrow \text{Al}^{3+}(aq) + 3\text{NO}_3^-(aq)$
 (D) $2\text{Ag}^+(aq) + \text{Ti}(s) \rightarrow 2\text{Ag}(s) + \text{Ti}^{2+}(aq)$

Use the following information to answer questions 46-50.





A sample of H_2S gas is placed in an evacuated, sealed container and heated until the following decomposition reaction occurs at 1000 K:



46. Which of the following represents the equilibrium constant for this reaction?

- (A) $K_c = \frac{[\text{H}_2]^2 [\text{S}_2]}{[\text{H}_2\text{S}]^2}$
 (B) $K_c = \frac{[\text{H}_2\text{S}]^2}{[\text{H}_2]^2 [\text{S}_2]}$
 (C) $K_c = \frac{2[\text{H}_2][\text{S}_2]}{2[\text{H}_2\text{S}]}$
 (D) $K_c = \frac{2[\text{H}_2\text{S}]}{2[\text{H}_2][\text{S}_2]}$

47. Which of the following graphs would best represent the change in concentration of the various species involved in the reaction over time?





48. Which option best describes what will immediately occur to the reaction rates if the pressure on the system is increased after it has reached equilibrium?
- (A) The rate of both the forward and reverse reactions will increase.
 (B) The rate of the forward reaction will increase while the rate of the reverse reaction decreases.
 (C) The rate of the forward reaction will decrease while the rate of the reverse reaction increases.
 (D) Neither the rate of the forward nor reverse reactions will change.
49. If, at a given point in the reaction, the value for the reaction quotient Q is determined to be 2.5×10^{-8} , which of the following is occurring?
- (A) The concentration of the reactant is decreasing while the concentration of the products is increasing.
 (B) The concentration of the reactant is increasing while the concentration of the products is decreasing.
 (C) The system has passed the equilibrium point, and the concentration of all species involved in the reaction will remain constant.
 (D) The concentrations of all species involved are changing at the same rate.

GO ON TO THE NEXT PAGE.

Section I


50. As the reaction progresses at a constant temperature of 1000 K, how does the value for the Gibbs free energy constant for the reaction change?
- (A) It stays constant.
 (B) It increases exponentially.
 (C) It increases linearly.
 (D) It decreases exponentially.
-
51. An unknown substance is found to have a high melting point. In addition, it is a poor conductor of electricity and does not dissolve in water. The substance most likely contains
- (A) ionic bonding
 (B) nonpolar covalent bonding
 (C) covalent network bonding
 (D) metallic bonding
52. Which of the following best explains why the ionization of atoms can occur during photoelectron spectroscopy, even though ionization is not a thermodynamically favored process?
- (A) It is an exothermic process due to the release of energy as an electron is liberated from the Coulombic attraction holding it to the nucleus.
 (B) The entropy of the system increases due to the separation of the electron from its atom.
 (C) Energy contained in the light can be used to overcome the Coulombic attraction between electrons and the nucleus.
 (D) The products of the ionization are at a lower energy state than the reactants.



The above diagrams show the decomposition of hydrogen peroxide in a sealed container in the presence of a catalyst. What is the overall order for the reaction?

- (A) Zero order
 (B) First order
 (C) Second order
 (D) Third order

54.



One of the resonance structures for the nitrite ion is shown above. What is the formal charge on each atom?

	O_x	N	O_y
(A)	-1	+1	-1
(B)	+1	-1	0
(C)	0	0	-1
(D)	-1	0	0


Use the following information to answer questions 55-57.

Atoms of four elements are examined: carbon, nitrogen, neon, and sulfur.

55. Atoms of which element would have the strongest magnetic moment?
- (A) Carbon
 (B) Nitrogen
 (C) Neon
 (D) Sulfur
56. Atoms of which element are most likely to form a structure with the formula XF_6 (where X is one of the four atoms)?
- (A) Carbon
 (B) Nitrogen
 (C) Neon
 (D) Sulfur
57. Which element would have a photoelectron spectra in which the peak representing electrons with the lowest ionization energy would be three times higher than all other peaks?
- (A) Carbon
 (B) Nitrogen
 (C) Neon
 (D) Sulfur

GO ON TO THE NEXT PAGE.

58. The diagram below supports which of the following conclusions about the reaction shown below?




- (A) There is an increase in entropy.
 (B) Mass is conserved in all chemical reactions.
 (C) The pressure increases after the reaction goes to completion.
 (D) The enthalpy value is positive.
59. $\text{NO}_2 + \text{O}_3 \rightarrow \text{NO}_3 + \text{O}_2$ slow
 $\text{NO}_3 + \text{NO}_2 \rightarrow \text{N}_2\text{O}_5$ fast

A proposed reaction mechanism for the reaction of nitrogen dioxide and ozone is detailed above. Which of the following is the rate law for the reaction?

- (A) $\text{Rate} = k[\text{NO}_2][\text{O}_3]$
 (B) $\text{Rate} = k[\text{NO}_3][\text{NO}_2]$
 (C) $\text{Rate} = k[\text{NO}_2]^2[\text{O}_3]$
 (D) $\text{Rate} = k[\text{NO}_3][\text{O}_2]$

- 60.



The concentrations of the reactants and products in the reaction represented by the above graph are found to be changing very slowly. Which of the following statements best describes the reaction given that the reaction is exergonic? ($\Delta G < 0$)

- (A) The reaction is under kinetic control.
 (B) The reaction has reached a state of equilibrium.
 (C) The reaction is highly exothermic in nature.
 (D) The addition of heat will increase the rate of reaction significantly.

END OF SECTION I

INFORMATION IN THE TABLE BELOW AND ON THE FOLLOWING PAGES MAY BE USEFUL IN ANSWERING THE QUESTIONS IN THIS SECTION OF THE EXAMINATION.

DO NOT DETACH FROM BOOK.

PERIODIC TABLE OF THE ELEMENTS

1	2																	3	4	5	6	7	8	9	10
H 1.008	He 4.00																	Li 6.94	Be 9.01	B 10.81	C 12.01	N 14.01	O 16.00	F 19.00	Ne 20.18
3	4																	11	12	13	14	15	16	17	18
Li 6.94	Be 9.01																	Na 22.99	Mg 24.30	Al 26.98	Si 28.09	P 30.97	S 32.06	Cl 35.45	Ar 39.95
19	20	21	22	23	24	25	26	27	28	29	30					31	32	33	34	35	36				
K 39.10	Ca 40.08	Sc 44.96	Ti 47.90	V 50.94	Cr 52.00	Mn 54.94	Fe 55.85	Co 58.93	Ni 58.69	Cu 63.55	Zn 65.39					Ga 69.72	Ge 72.59	As 74.92	Se 78.96	Br 79.90	Kr 83.80				
37	38	39	40	41	42	43	44	45	46	47	48					49	50	51	52	53	54				
Rb 85.47	Sr 87.62	Y 88.91	Zr 91.22	Nb 92.91	Mo 95.94	Tc (98)	Ru 101.1	Rh 102.91	Pd 106.42	Ag 107.87	Cd 112.41					In 114.82	Sn 118.71	Sb 121.75	Te 127.60	I 126.91	Xe 131.29				
55	56	57	72	73	74	75	76	77	78	79	80					81	82	83	84	85	86				
Cs 132.91	Ba 137.33	*La 138.91	Hf 178.49	Ta 180.95	W 183.85	Re 186.21	Os 190.2	Ir 192.2	Pt 195.08	Au 196.97	Hg 200.59					Tl 204.38	Pb 207.2	Bi 208.98	Po (209)	At (210)	Rn (222)				
87	88	89	104	105	106	107	108	109	110	111					112	113	114	115	116	117	118				
Fr (223)	Ra 226.02	†Ac 227.03	Rf (261)	Db (262)	Sg (266)	Bh (264)	Hs (277)	Mt (268)	Ds (271)	Rg (272)					119	120	121	122	123	124	125	126			
				58	59	60	61	62	63	64	65	66	67	68	69	70	71								
				Ce 140.12	Pr 140.91	Nd 144.24	Pm (145)	Sm 150.4	Eu 151.97	Gd 157.25	Tb 158.93	Dy 162.50	Ho 164.93	Er 167.26	Tm 168.93	Yb 173.04	Lu 174.97								
				90	91	92	93	94	95	96	97	98	99	100	101	102	103								
				Th 232.04	Pa 231.04	U 238.03	Np (237)	Pu (244)	Am (243)	Cm (247)	Bk (247)	Cf (251)	Es (252)	Fm (257)	Md (258)	No (259)	Lr (262)								

*Lanthanide Series

† Actinide Series

GO ON TO THE NEXT PAGE.

ADVANCED PLACEMENT CHEMISTRY EQUATIONS AND CONSTANTS

Throughout the test the following symbols have the definitions specified unless otherwise noted.

L, mL = liter(s), milliliter(s)
 g = gram(s)
 nm = nanometer(s)
 atm = atmosphere(s)

mm Hg = millimeters of mercury
 J, kJ = joule(s), kilojoule(s)
 V = volt(s)
 mol = mole(s)

ATOMIC STRUCTURE

$$E = h\nu$$

$$c = \lambda\nu$$

E = energy
 ν = frequency
 λ = wavelength

Planck's constant, $h = 6.626 \times 10^{-34}$ J s

Speed of light, $c = 2.998 \times 10^8$ m s⁻¹

Avogadro's number = 6.022×10^{23} mol⁻¹

Electron charge, $e = -1.602 \times 10^{-19}$ coulomb

EQUILIBRIUM

$$K_c = \frac{[C]^c[D]^d}{[A]^a[B]^b}, \text{ where } aA + bB \rightleftharpoons cC + dD$$

$$K_p = \frac{(P_C)^c(P_D)^d}{(P_A)^a(P_B)^b}$$

$$K_a = \frac{[H^+][A^-]}{[HA]}$$

$$K_b = \frac{[OH^-][HB^+]}{[B]}$$

$$K_w = [H^+][OH^-] = 1.0 \times 10^{-14} \text{ at } 25^\circ\text{C}$$

$$= K_a \times K_b$$

$$\text{pH} = -\log[H^+], \text{ pOH} = -\log[OH^-]$$

$$14 = \text{pH} + \text{pOH}$$

$$\text{pH} = \text{p}K_a + \log \frac{[A^-]}{[HA]}$$

$$\text{p}K_a = -\log K_a, \text{ p}K_b = -\log K_b$$

Equilibrium Constants

K_c (molar concentrations)

K_p (gas pressures)

K_a (weak acid)

K_b (weak base)

K_w (water)

KINETICS

$$\ln[A]_t - \ln[A]_0 = -kt$$

$$\frac{1}{[A]_t} - \frac{1}{[A]_0} = kt$$

$$t_{1/2} = \frac{0.693}{k}$$

k = rate constant

t = time

$t_{1/2}$ = half-life

GO ON TO THE NEXT PAGE.

GASES, LIQUIDS, AND SOLUTIONS

$$PV = nRT$$

$$P_A = P_{\text{total}} \times X_A, \text{ where } X_A = \frac{\text{moles A}}{\text{total moles}}$$

$$P_{\text{total}} = P_A + P_B + P_C + \dots$$

$$n = \frac{m}{M}$$

$$K = ^\circ\text{C} + 273$$

$$D = \frac{m}{V}$$

$$KE \text{ per molecule} = \frac{1}{2}mv^2$$

Molarity, M = moles of solute per liter of solution

$$A = abc$$

P = pressure

V = volume

T = temperature

n = number of moles

m = mass

M = molar mass

D = density

KE = kinetic energy

v = velocity

A = absorbance

a = molar absorptivity

b = path length

c = concentration

Gas constant, $R = 8.314 \text{ J mol}^{-1} \text{ K}^{-1}$

$$= 0.08206 \text{ L atm mol}^{-1} \text{ K}^{-1}$$

$$= 62.36 \text{ L torr mol}^{-1} \text{ K}^{-1}$$

$$1 \text{ atm} = 760 \text{ mm Hg}$$

$$= 760 \text{ torr}$$

STP = 273.15 K and 1.0 atm

Ideal gas at STP = 22.4 L mol⁻¹

THERMOCHEMISTRY/ ELECTROCHEMISTRY

$$q = mc\Delta T$$

$$\Delta S^\circ = \sum S^\circ \text{ products} - \sum S^\circ \text{ reactants}$$

$$\Delta H^\circ = \sum \Delta H_f^\circ \text{ products} - \sum \Delta H_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \sum \Delta G_f^\circ \text{ products} - \sum \Delta G_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

$$= -RT \ln K$$

$$= -nFE^\circ$$

$$I = \frac{q}{t}$$

q = heat

m = mass

c = specific heat capacity

T = temperature

S° = standard entropy

H° = standard enthalpy

G° = standard free energy

n = number of moles

E° = standard reduction potential

I = current (amperes)

q = charge (coulombs)

t = time (seconds)

Faraday's constant, $F = 96,485$ coulombs per mole of electrons

$$1 \text{ volt} = \frac{1 \text{ joule}}{1 \text{ coulomb}}$$

GO ON TO THE NEXT PAGE.

CHEMISTRY

Section II

7 Questions

(Total time—1 hour and 45 minutes)

YOU MAY USE YOUR CALCULATOR FOR THIS SECTION.

Directions: Questions 1-3 are long free-response questions that require about 23 minutes each to answer and are worth 10 points each. Questions 4-7 are short free-response questions that require about 9 minutes each to answer and are worth 4 points each.

Write your response in the space provided following each question. Examples and equations may be included in your responses where appropriate. For calculations, clearly show the method used and the steps involved in arriving at your answers. You must show your work to receive credit for your answer. Pay attention to significant figures.

1. A stock solution of 0.100 *M* cobalt (II) chloride is used to create several solutions, indicated in the data table below:

Sample	Volume CoCl ₂ (mL)	Volume H ₂ O (mL)
1	20.00	0
2	15.00	5.00
3	10.00	10.00
4	5.00	15.00

- (a) In order to achieve the degree of accuracy shown in the table above, select which of the following pieces of laboratory equipment could be used when measuring out the CoCl₂:

150-mL beaker
50-mL buret

400-mL beaker
50-mL graduated cylinder

250-mL Erlenmeyer flask
100-mL graduated cylinder


- (b) Calculate the concentration of the CoCl₂ in each sample.

The solutions are then placed in cuvettes before being inserted into a spectrophotometer calibrated to 560 nm and their values are measured, yielding the data below:

Sample	Absorbance
1	0.485
2	0.364
3	0.243
4	0.121


- (c) If gloves are not worn when handling the cuvettes, how might this affect the absorbance values gathered?
- (d) If the path length of the cuvette is 1.00 cm, what is the molar absorptivity value for CoCl₂ at 560 nm?
- (e) On the axes on the next page, plot a graph of absorbance vs. concentration. The y-axis scale is set, and be sure to scale the x-axis appropriately

GO ON TO THE NEXT PAGE.



- (f) What would the absorbance values be for CoCl_2 solutions at the following concentrations?
- (i) 0.067 M
 - (ii) 0.180 M

GO ON TO THE NEXT PAGE.



2. A sample of liquid butane (C_4H_{10}) in a pressurized lighter is set up directly beneath an aluminum can, as shown in the diagram above. The can contains 100.0 mL of water, and when the butane is ignited the temperature of the water inside the can increases from 25.0°C to 82.3°C. The total mass of butane ignited is found to be 0.51 g, the specific heat of water is 4.18 J/g·°C, and the density of water is 1.00 g/mL.
- Write the balanced chemical equation for the combustion of one mole of butane in air.
 - How much heat did the water gain?
 - What is the experimentally determined heat of combustion for butane based on this experiment? Your answer should be in kJ/mol.
 - Given butane's density of 0.573 g/mL at 25°C, calculate how much heat would be emitted if 5.00 mL of it were combusted at that temperature.
 - The overall combustion of butane is an exothermic reaction. Explain why this is, in terms of bond energies.
 - One of the major sources of error in this experiment comes from the heat that is absorbed by the air. Why, then, might it not be a good idea to perform this experiment inside a sealed container to prevent the heat from leaving the system?

GO ON TO THE NEXT PAGE.

Section II



The data below was gathered for the decomposition of N_2O_5 at 310 K via the equation above.

Time (s)	$[\text{N}_2\text{O}_5] (M)$
0	0.250
500.	0.190
1000.	0.145
2000.	0.085


- (a) How does the rate of appearance of NO_2 compare to the rate of disappearance of N_2O_5 ? Justify your answer.
- (b) The reaction is determined to be first order overall. On the axes below, create a graph of some function of concentration vs. time that will produce a straight line. Label and scale your axes appropriately.




- (c) (i) What is the rate constant for this reaction? Include units.
 (ii) What would the concentration of N_2O_5 be at $t = 1500$ s?
 (iii) What is the half-life of N_2O_5 ?
- (d) Would the addition of a catalyst increase, decrease, or have no effect on the following variables? Justify your answers.
 (i) Rate of disappearance of N_2O_5
 (ii) Magnitude of the rate constant
 (iii) Half-life of N_2O_5

GO ON TO THE NEXT PAGE.


4. Consider the Lewis structures for the following four molecules:




n-Butylamine



Propanal




Pentane



Methanol

- (a) All of the substances are liquids at room temperature. Organize them from high to low in terms of boiling points, clearly differentiating between the intermolecular forces in each substance.
- (b) On the methanol diagram reproduced below, draw the locations of all dipoles.




Methanol

- (c) n-Butylamine is found to have the lowest vapor pressure at room temperature out of the four liquids. Justify this observation in terms of intermolecular forces.

GO ON TO THE NEXT PAGE.

Section II

5. Current is run through an aqueous solution of nickel (II) fluoride, and a gas is evolved at the right-hand electrode, as indicated by the diagram below:



The standard reduction potential for several reactions is given in the following table:

Half-cell	E°_{red}
$\text{F}_2(g) + 2e^- \rightarrow 2\text{F}^-$	+2.87 V
$\text{O}_2(g) + 4\text{H}^+ + 4e^- \rightarrow 2\text{H}_2\text{O}(l)$	+1.23 V
$\text{Ni}^{2+} + 2e^- \rightarrow \text{Ni}(s)$	-0.25 V
$2\text{H}_2\text{O}(l) + 2e^- \rightarrow \text{H}_2(g) + 2\text{OH}^-$	-0.83 V

- (a) Determine which half-reaction is occurring at each electrode:
- Oxidation
 - Reduction
- (b) (i) Calculate the standard cell potential of the cell.
 (ii) Calculate the Gibbs free energy value of the cell at standard conditions.
- (c) Which electrode in the diagram (A or B) is the cathode, and which is the anode? Justify your answers.

GO ON TO THE NEXT PAGE.

6. Aniline, $C_6H_5NH_2$, is a weak base with $K_b = 3.8 \times 10^{-10}$.
- Write out the reaction that occurs when aniline reacts with water.
 - What is the concentration of each species at equilibrium in a solution of $0.25 M C_6H_5NH_2$?
 - What is the pH value for the solution in (i)?
7. A rigid, sealed 12.00 L container is filled with 10.00 g each of three different gases: CO_2 , NO , and NH_3 . The temperature of the gases is held constant $35.0^\circ C$. Assume ideal behavior for all gases.
- What is the mole fraction of each gas?
 - What is the partial pressure of each gas?
 - Out of the three gases, molecules of which gas will have the highest velocity? Why?
 - Name one circumstance in which the gases might deviate from ideal behavior, and clearly explain the reason for the deviation.

STOP

END OF EXAM

Completely darken bubbles with a No. 2 pencil. If you make a mistake, be sure to erase mark completely. Erase all stray marks.

1. YOUR NAME: _____
(Print) Last First M.I.

SIGNATURE: _____ DATE: ____/____/____

HOME ADDRESS: _____
(Print) Number and Street

 City State Zip Code

PHONE NO.: _____

5. YOUR NAME

First 4 letters of last name				FIRST INIT	MID INIT
(A)	(A)	(A)	(A)	(A)	(A)
(B)	(B)	(B)	(B)	(B)	(B)
(C)	(C)	(C)	(C)	(C)	(C)
(D)	(D)	(D)	(D)	(D)	(D)
(E)	(E)	(E)	(E)	(E)	(E)
(F)	(F)	(F)	(F)	(F)	(F)
(G)	(G)	(G)	(G)	(G)	(G)
(H)	(H)	(H)	(H)	(H)	(H)
(I)	(I)	(I)	(I)	(I)	(I)
(J)	(J)	(J)	(J)	(J)	(J)
(K)	(K)	(K)	(K)	(K)	(K)
(L)	(L)	(L)	(L)	(L)	(L)
(M)	(M)	(M)	(M)	(M)	(M)
(N)	(N)	(N)	(N)	(N)	(N)
(O)	(O)	(O)	(O)	(O)	(O)
(P)	(P)	(P)	(P)	(P)	(P)
(Q)	(Q)	(Q)	(Q)	(Q)	(Q)
(R)	(R)	(R)	(R)	(R)	(R)
(S)	(S)	(S)	(S)	(S)	(S)
(T)	(T)	(T)	(T)	(T)	(T)
(U)	(U)	(U)	(U)	(U)	(U)
(V)	(V)	(V)	(V)	(V)	(V)
(W)	(W)	(W)	(W)	(W)	(W)
(X)	(X)	(X)	(X)	(X)	(X)
(Y)	(Y)	(Y)	(Y)	(Y)	(Y)
(Z)	(Z)	(Z)	(Z)	(Z)	(Z)

IMPORTANT: Please fill in these boxes exactly as shown on the back cover of your test book.

2. TEST FORM

3. TEST CODE				4. REGISTRATION NUMBER																
(0)	(A)	(J)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0)
(1)	(B)	(K)	(1)	(1)	(1)	(1)	(1)	(1)	(1)	(1)	(1)	(1)	(1)	(1)	(1)	(1)	(1)	(1)	(1)	(1)
(2)	(C)	(L)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)	(2)
(3)	(D)	(M)	(3)	(3)	(3)	(3)	(3)	(3)	(3)	(3)	(3)	(3)	(3)	(3)	(3)	(3)	(3)	(3)	(3)	(3)
(4)	(E)	(N)	(4)	(4)	(4)	(4)	(4)	(4)	(4)	(4)	(4)	(4)	(4)	(4)	(4)	(4)	(4)	(4)	(4)	(4)
(5)	(F)	(O)	(5)	(5)	(5)	(5)	(5)	(5)	(5)	(5)	(5)	(5)	(5)	(5)	(5)	(5)	(5)	(5)	(5)	(5)
(6)	(G)	(P)	(6)	(6)	(6)	(6)	(6)	(6)	(6)	(6)	(6)	(6)	(6)	(6)	(6)	(6)	(6)	(6)	(6)	(6)
(7)	(H)	(Q)	(7)	(7)	(7)	(7)	(7)	(7)	(7)	(7)	(7)	(7)	(7)	(7)	(7)	(7)	(7)	(7)	(7)	(7)
(8)	(I)	(R)	(8)	(8)	(8)	(8)	(8)	(8)	(8)	(8)	(8)	(8)	(8)	(8)	(8)	(8)	(8)	(8)	(8)	(8)
(9)			(9)	(9)	(9)	(9)	(9)	(9)	(9)	(9)	(9)	(9)	(9)	(9)	(9)	(9)	(9)	(9)	(9)	(9)

6. DATE OF BIRTH

Month	Day		Year	
() JAN				
() FEB	(0)	(0)	(0)	(0)
() MAR	(1)	(1)	(1)	(1)
() APR	(2)	(2)	(2)	(2)
() MAY	(3)	(3)	(3)	(3)
() JUN		(4)	(4)	(4)
() JUL		(5)	(5)	(5)
() AUG		(6)	(6)	(6)
() SEP		(7)	(7)	(7)
() OCT		(8)	(8)	(8)
() NOV		(9)	(9)	(9)
() DEC				

7. GENDER

() MALE

() FEMALE



1. (A) (B) (C) (D)	21. (A) (B) (C) (D)	41. (A) (B) (C) (D)
2. (A) (B) (C) (D)	22. (A) (B) (C) (D)	42. (A) (B) (C) (D)
3. (A) (B) (C) (D)	23. (A) (B) (C) (D)	43. (A) (B) (C) (D)
4. (A) (B) (C) (D)	24. (A) (B) (C) (D)	44. (A) (B) (C) (D)
5. (A) (B) (C) (D)	25. (A) (B) (C) (D)	45. (A) (B) (C) (D)
6. (A) (B) (C) (D)	26. (A) (B) (C) (D)	46. (A) (B) (C) (D)
7. (A) (B) (C) (D)	27. (A) (B) (C) (D)	47. (A) (B) (C) (D)
8. (A) (B) (C) (D)	28. (A) (B) (C) (D)	48. (A) (B) (C) (D)
9. (A) (B) (C) (D)	29. (A) (B) (C) (D)	49. (A) (B) (C) (D)
10. (A) (B) (C) (D)	30. (A) (B) (C) (D)	50. (A) (B) (C) (D)
11. (A) (B) (C) (D)	31. (A) (B) (C) (D)	51. (A) (B) (C) (D)
12. (A) (B) (C) (D)	32. (A) (B) (C) (D)	52. (A) (B) (C) (D)
13. (A) (B) (C) (D)	33. (A) (B) (C) (D)	53. (A) (B) (C) (D)
14. (A) (B) (C) (D)	34. (A) (B) (C) (D)	54. (A) (B) (C) (D)
15. (A) (B) (C) (D)	35. (A) (B) (C) (D)	55. (A) (B) (C) (D)
16. (A) (B) (C) (D)	36. (A) (B) (C) (D)	56. (A) (B) (C) (D)
17. (A) (B) (C) (D)	37. (A) (B) (C) (D)	57. (A) (B) (C) (D)
18. (A) (B) (C) (D)	38. (A) (B) (C) (D)	58. (A) (B) (C) (D)
19. (A) (B) (C) (D)	39. (A) (B) (C) (D)	59. (A) (B) (C) (D)
20. (A) (B) (C) (D)	40. (A) (B) (C) (D)	60. (A) (B) (C) (D)