## CHEMISTRY SUBJECT TEST

### MATERIAL IN THE FOLLOWING TABLE MAY BE USEFUL IN ANSWERING THE QUESTIONS IN THIS EXAMINATION.

Г

DO NOT DETACH FROM BOOK.

|          | 2 | He | 4.0026 | 10 | Ne | 20.179 | 18 | $\mathbf{Ar}$ | 39.948 | 36 | Kr | 83.80  | 54 | Хе            | 131.29 | 86        | Rn  | (222)  |     |           |        |    |               |
|----------|---|----|--------|----|----|--------|----|---------------|--------|----|----|--------|----|---------------|--------|-----------|-----|--------|-----|-----------|--------|----|---------------|
|          |   |    |        | 6  | F  | 19.00  | 17 | CI            | 35.453 | 35 | Br | 79.90  | 53 | I             | 126.91 | 85        | At  | (210)  |     |           |        | 71 | Lu            |
| S        |   |    |        | 8  | 0  | 16.00  | 16 | s             | 32.06  | 34 | Se | 78.96  | 52 | Te            | 127.60 | 84        | Po  | (209)  |     |           |        | 70 | Yb            |
| Z        |   |    |        | 7  | Z  | 14.007 | 15 | Ρ             | 30.974 | 33 | As | 74.92  | 51 | $\mathbf{Sb}$ | 121.75 | 83        | Bi  | 208.98 |     |           |        | 69 | Tm            |
| M        |   |    |        | 9  | C  | 12.011 | 14 | Si            | 28.09  | 32 | Ge | 72.59  | 50 | Sn            | 118.71 | 82        | Рb  | 207.2  |     | yet named |        | 68 | Er            |
|          |   |    |        | ъ  | B  | 10.811 | 13 | AI            | 26.98  | 31 | Ga | 69.72  | 49 | In            | 114.82 | 81        | П   | 204.38 |     | § Not     |        | 67 | Ηo            |
| ш        |   |    |        |    |    |        |    |               |        | 30 | Zn | 65.39  | 48 | Cd            | 112.41 | 80        | Hg  | 200.59 | 112 | so        | (277)  | 99 | Dy            |
| 폰        |   |    |        |    |    |        |    |               |        | 29 | Cu | 63.55  | 47 | Ag            | 107.87 | 79        | Νu  | 196.97 | 111 | Rg        | (272)  | 65 | τb            |
| 0Е       |   |    |        |    |    |        |    |               |        | 28 | Ni | 58.69  | 46 | Ρd            | 106.42 | 78        | Pt  | 195.08 | 110 | $D_{S}$   | (271)  | 64 | Gd            |
| Ľ        |   |    |        |    |    |        |    |               |        | 27 | Co | 58.93  | 45 | Rh            | 102.91 | <i>LL</i> | Ir  | 192.2  | 109 | Mt        | (268)  | 63 | Eu            |
| AB       |   |    |        |    |    |        |    |               |        | 26 | Fe | 55.85  | 44 | Ru            | 101.1  | 92        | Os  | 190.2  | 108 | $H_{S}$   | (277)  | 62 | Sm            |
| C        |   |    |        |    |    |        |    |               |        | 25 | Mn | 54.938 | 43 | Tc            | (86)   | 75        | Re  | 186.21 | 107 | Bh        | (264)  | 61 | Pm            |
| IOO      |   |    |        |    |    |        |    |               |        | 24 | C  | 52.00  | 42 | Mo            | 95.94  | 74        | Μ   | 183.85 | 106 | Sg        | (266)  | 60 | ΡN            |
| BI       |   |    |        |    |    |        |    |               |        | 23 | Λ  | 50.94  | 41 | ЧN            | 92.91  | 73        | Та  | 180.95 | 105 | Db        | (262)  | 59 | $\mathbf{Pr}$ |
| <b>L</b> |   |    |        |    |    |        |    |               |        | 22 | Τi | 47.90  | 40 | Zr            | 91.22  | 72        | Hf  | 178.49 | 104 | Rf        | (261)  | 58 | Ce            |
|          |   |    |        |    |    |        |    |               |        | 21 | Sc | 44.96  | 39 | Υ             | 88.91  | 22        | *La | 138.91 | 68  | ⁺Αc       | 227.03 |    | ries          |
|          |   |    |        | 4  | Be | 9.012  | 12 | Mg            | 24.30  | 20 | Ca | 40.48  | 38 | Sr            | 87.62  | 56        | Ba  | 137.33 | 88  | Ra        | 226.02 |    | hanide Se     |
|          | - | Η  | 1.0079 | ю  | Li | 6.941  | 11 | Na            | 22.99  | 19 | Х  | 39.10  | 37 | Rb            | 85.47  | 55        | Cs  | 132.91 | 87  | Fr        | (223)  |    | *Lant         |

**GO ON TO THE NEXT PAGE** 

174.97

173.04

168.93

167.26

164.93 Нo

162.50

158.93

157.25

151.97

150.4

(145)Pm

144.24

140.91

140.12

95

94 Pu

93

92 U

91 Pa

90 Th

<sup>†</sup>Actinide Series

Dy

103

102

101

100 Fm

66  $\mathbf{E}_{\mathbf{S}}$ 

98 Ç (262)

(259) °N

(258)

(257)

(252)

(251)

(247)

(247)

(243)Am

(244)

237.05

238.03

231.04

232.04

Np

Bk 97

Cm 96

Ľ

Мd

### **CHEMISTRY SUBJECT TEST 1**

<u>Note:</u> For all questions involving solutions and/or chemical equations, assume that the system is in pure water unless otherwise stated.

### Part A

**Directions:** Each set of lettered choices below refers to the numbered statements or questions immediately following it. Select the one lettered choice that best fits each statement or answers each question, and then fill in the corresponding oval on the answer sheet. A choice may be used once, more than once, or not at all in each set.

Questions 1–4 refer to the following.

- (A) Thermometer
- (B) Conductivity tester
- (C) Salt bridge
- (D) Buret
- (E) Graduated cylinder
- 1. May be used in combination with a calorimeter to compare the specific heats of two substances
- 2. Is used to measure the volume of a solid by water displacement
- 3. Useful for adding small quantities of acid into a base
- 4. Completes the circuit of an electrochemical cell

Questions 5–9 refer to the following.

- (A) Nucleic acids
- (B) Proteins
- (C) Carbohydrates
- (D) Lipids
- (E) Electrolytes
- 5. Always amphoteric in nature
- 6. Found as both straight-chained and branched polymers
- 7. Deoxyribose in DNA nucleotides belongs to this family of biologically important molecules
- 8. Always ionic in nature
- 9. Tend not to be water soluble, and aggregate into droplets or molecular bilayers



### Questions 10–13 refer to the following.

- (A)  $Ag^+ + Br^- \rightarrow AgBr$
- (B)  ${}^{14}_{6}C \rightarrow {}^{14}_{7}N + {}^{0}_{-1}e$
- (C)  $^{234}_{92}U \rightarrow ^{230}_{90}Th + ^{4}_{2}He$
- (D)  ${}^{30}_{15}P \rightarrow {}^{30}_{14}Si + {}^{0}_{1}e$
- (E)  $2HgO \rightarrow 2Hg + O_2$
- 10. Represents the decomposition of a compound into its constituent elements
- 11. Represents alpha decay
- 12. Represents an oxidation-reduction reaction
- 13. Causes the neutron-to-proton ratio in a nucleus to be lowered

### Questions 14–16 refer to the following.



- 14. Is the activation energy of the reverse reaction
- 15. Is the enthalpy change of the forward reaction
- 16. Represents energy of the activated complex



### Questions 17–20 refer to the following.

- (A) Hydrogen bonding
- (B) Ionic bonding
- (C) Metallic bonding
- (D) Nonpolar covalent bonding
- (E) Polar covalent bonding
- 17. Holds a sample of barium iodide, BaI<sub>2</sub>, together
- 18. Allows solids to conduct electricity
- 19. Attracts atoms of hydrogen to each other in an  $H_2$  molecule
- 20. Responsible for relatively low vapor pressure of water

### Questions 21–23 refer to the following.

- (A) Iron(III) chloride,  $FeCl_3(s)$
- (B) Iodine,  $I_2(s)$
- (C) Sodium hydroxide, NaOH(s)
- (D) Sucrose,  $C_{12}H_{22}O_{11}(s)$
- (E) Graphite, C(s)
- 21. Gives off a purplish vapor as it sublimes
- 22. Can conduct electricity in the solid state
- 23. Its dissolution in water is highly exothermic



### PLEASE GO TO THE SPECIAL SECTION LABELED CHEMISTRY AT THE LOWER RIGHT-HAND CORNER OF THE ANSWER SHEET YOU ARE WORKING ON AND ANSWER QUESTIONS 101–116 ACCORDING TO THE FOLLOWING DIRECTIONS.

#### Part B

**Directions:** Each question below consists of two statements, I in the left-hand column and II in the right-hand column. For each question, determine whether statement I is true or false <u>and</u> whether statement II is true or false, and fill in the corresponding T or F ovals on your answer sheet. <u>Fill in oval CE only if statement II is a correct explanation of statement I.</u>

| EXAMPLES:  |         |  |
|--|---------|--|
| Ī  |         | Ш  |
| <b>EX 1.</b> $H_2$ <b>SO</b> <sub>4</sub> is a strong acid | BECAUSE | $H_2 SO_4$ contains sulfur.  |
| EX 2. An atom of oxygen is electrically neutral            | BECAUSE | an oxygen atom contains an equal<br>number of protons and electrons. |
| SAMPLE ANSWERS   |         |  |
| EX 1<br>EX 2   | I II CE |  |

|      | Ī  |         | <u>II</u>  |
|------|--|---------|--|
| 101. | Carbon is a nonmetal   | BECAUSE | carbon atoms can bond with each other.   |
| 102. | Two isotopes of the same element have the same mass number       | BECAUSE | isotopes have the same number of protons.  |
| 103. | The density of a sample of water is doubled by doubling its mass | BECAUSE | compared to a gas, the molecules in a liquid are relatively far apart.                             |
| 104. | Sodium and cesium exhibit similar chemical properties            | BECAUSE | their atoms have the same number of valence electrons.   |
| 105. | An endothermic reaction can be spontaneous                       | BECAUSE | both enthalpy and entropy changes<br>affect the value of a reaction's Gibbs<br>free energy change. |
| 106. | The 4 <i>s</i> orbital fills before the $3d$ orbitals            | BECAUSE | subshells fill in order from lower<br>to higher energy.  |
| 107. | Calcium acts as a reducing agent when it reacts with bromine     | BECAUSE | mass is conserved in a chemical reaction. <b>GO ON TO THE NEXT PAGE</b>                            |

### Ī

### II

| 108. | If an acid is added to pure water, it increases the water's pH                            | BECAUSE | adding an acid to water raises the<br>hydrogen ion concentration in the<br>water.             |
|------|---|---------|---|
| 109. | Covalent bonds must be broken for a liquid to boil  | BECAUSE | heat must be released for a liquid to change into a gas.                                      |
| 110. | Alpha particles can be detected using a Geiger counter                                    | BECAUSE | all radioactive elements are highly chemically reactive.                                      |
| 111. | As ice absorbs heat and begins to<br>melt, its temperature remains<br>constant            | BECAUSE | the absorbed heat is consumed by the breaking of intermolecular interactions.                 |
| 112. | When a solute is added to pure<br>water, the vapor pressure of the<br>water will decrease | BECAUSE | all solutes dissociate into positive and negative ions.                                       |
| 113. | The rate of a reaction is accelerated by increasing temperature                           | BECAUSE | a large equilibrium constant favors the formation of product.                                 |
| 114. | Hydrofluoric acid, $HF(aq)$ , is a weaker electrolyte than hydrochloric acid, $HCl(aq)$ , | BECAUSE | fluorine has a lower electronegativity than chlorine.   |
| 115. | A nonpolar molecule can have polar bonds  | BECAUSE | polar bonds can be symmetrically<br>arranged in a molecule so that there are<br>no net poles. |
| 116. | The electrolysis of potassium iodide,<br>KI, produces electrical energy                   | BECAUSE | electrons flow from the anode to the cathode.   |

RETURN TO THE SECTION OF YOUR ANSWER SHEET YOU STARTED FOR **CHEMISTRY** AND ANSWER QUESTIONS 24–69.



### Part C

**Directions:** Each of the questions or incomplete statements below is followed by five suggested answers or completions. Select the one that is best in each case and then fill in the corresponding oval on the answer sheet.

- 24. What is the number of protons and neutrons in an atom with mass number 89 and atomic number 39 ?
  - (A) 50 protons and 50 neutrons
  - (B) 50 protons and 39 neutrons
  - (C) 39 protons and 89 neutrons
  - (D) 39 protons and 50 neutrons
  - (E) 39 protons and 39 neutrons
  - $\dots C_{\underline{a}}H_{\underline{10}}(g) + \dots O_{\underline{a}}(g) \rightarrow \dots CO_{\underline{a}}(g) + \dots H_{\underline{a}}O(l)$
- 25. When the above equation is balanced using the lowest whole-number terms, the coefficient of  $CO_2$  is
  - (A) 2
  - (B) 4
  - (C) 8
  - (D) 10
  - (E) 13
- 26. Which of the following is closest in mass to a proton?
  - (A) Alpha particle
  - (B) Positron
  - (C) Neutron
  - (D) Electron
  - (E) Hydrogen molecule
- 27. What is the approximate percentage composition by mass of the element oxygen in the compound  $HClO_{a}$ ?
  - (A) 16%
  - (B) 32%
  - (C) 50%
  - (D) 64%
  - (E) 75%

- 28. If two atoms that differ in electronegativity combine by chemical reaction and share electrons, the bond that joins them will be
  - (A) metallic
  - (B) ionic
  - (C) a hydrogen bond
  - (D) nonpolar covalent
  - (E) polar covalent
- 29. When the temperature of a 20-gram sample of water is increased from 10°C to 30°C, the heat transferred to the water is
  - (A) 600 calories
  - (B) 400 calories
  - (C) 200 calories
  - (D) 30 calories
  - (E) 20 calories
- 30. What is the oxidation state of chromium, Cr, in the compound potassium dichromate, K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub>?
  - (A) +1
  - (B) +2
  - (C) +3
  - (D) +6
  - (E) +12
- 31. An aqueous solution with pH 5 at 25°C has a hydroxide ion (OH<sup>-</sup>) concentration of
  - (A)  $1 \times 10^{-11}$  molar
  - (B)  $1 \times 10^{-9}$  molar
  - (C)  $1 \times 10^{-7}$  molar
  - (D)  $1 \times 10^{-5}$  molar
  - (E)  $1 \times 10^{-3}$  molar



 $2\mathrm{H}_{2}\mathrm{O}(g) \rightarrow 2\mathrm{H}_{2}(g) + \mathrm{O}_{2}(g)$ 

- 32. The volume of water vapor required to produce 44.8 liters of oxygen by the above reaction is
  - (A) 11.2 liters
  - (B) 22.4 liters
  - (C) 44.8 liters
  - (D) 89.6 liters
  - (E) 100.0 liters
- 33. When 190 grams of MgCl<sub>2</sub> are dissolved in water and the resulting solution is 500 milliliters in volume, what is the molar concentration of MgCl<sub>2</sub> in the solution?
  - (A) 2.0 *M*
  - (B) 4.0 *M*
  - (C) 8.0 *M*
  - (D) 12.0 *M*
  - (E) 16.0 *M*
- 34. When a fixed amount of gas has its Kelvin temperature doubled and its pressure doubled, the new volume of the gas is
  - (A) four times greater than its original volume
  - (B) twice its original volume
  - (C) unchanged
  - (D) one-half its original volume
  - (E) one-fourth its original volume
- 35. In 12.4 hours, a 100 gram sample of an element decays so that its mass is 25 grams. What is the approximate half-life of this radioactive substance?
  - (A) 1.6 hours
  - (B) 3.1 hours
  - (C) 6.2 hours
  - (D) 24.8 hours
  - (E) 49.6 hours

- 36. In the equation Q  $\rightarrow {}^{4}_{2}He + {}^{216}_{85}At$ , the species represented by Q is
  - (A)  $^{220}_{87}$  Fr
  - (B)  $^{212}_{83}$ Bi
  - (C)  $^{220}_{87}$ At
  - (D)  $^{212}_{83}$ Fr
  - (E) <sup>216</sup><sub>85</sub>Bi
- 37. A compound with a molecular weight of 56 amu has an empirical formula of  $CH_2$ . What is its molecular formula?
  - (A)  $C_2H_2$
  - (B)  $C_2H_4$
  - $(C) C_4 H_8$
  - (D)  $C_4 H_{10}$
  - (E)  $C_6 H_{12}$
- 38. The change in heat energy for a reaction is best expressed as a change in
  - (A) enthalpy
  - (B) absolute temperature
  - (C) specific heat
  - (D) entropy
  - (E) kinetic energy

 $\dots \operatorname{NF}_3(g) + \dots + \operatorname{H}_2\operatorname{O}(g) \to \dots + \operatorname{HF}(g) + \dots \operatorname{NO}(g) + \dots$  $\dots \operatorname{NO}_2(g)$ 

- 39. When the equation for the reaction above is balanced, how many moles of NF<sub>3</sub> would be required to react completely with 6 moles of H<sub>2</sub>O ?
  - (A) 0.5 mole
  - (B) 1 mole
  - (C) 2 moles
  - (D) 3 moles
  - (E) 4 moles



- 40. Which characteristic is associated with bases?
  - (A) React with metal to produce hydrogen gas
  - (B) Donate an unshared electron pair
  - (C) Always contain the hydroxide ion in their structure
  - (D) Taste sour
  - (E) Formed by the reaction of a nonmetal oxide and water
- 41. An element has the following properties: shiny, brittle, poor electrical conductivity, and high melting point. This element can be best classified as a(n)
  - (A) alkali metal
  - (B) halogen
  - (C) metalloid
  - (D) transition metal
  - (E) noble gas
- 42. Which of the following forward processes produces a decrease in entropy?
  - I.  $H_2O(g) \rightarrow H_2O(l)$
  - II.  $\operatorname{Fe}^{2+}(aq) + \operatorname{S}^{2-}(aq) \to \operatorname{FeS}(s)$
  - III.  $2SO_3(g) \rightleftharpoons 2SO_2(g) + O_2(g)$
  - (A) I only
  - (B) III only
  - (C) I and II only
  - $(D) \ \ II \ and \ III \ only$
  - (E) I, II, and III
- 43. Which of the following will raise the boiling point of a sample of water?
  - (A) Heat the water
  - (B) Mix gasoline into the water
  - (C) Bring the water sample to a higher altitude
  - (D) Place the water sample on a magnetic stirrer
  - (E) Dissolve table sugar into the water

- 44. Elements H and J lie in the same period. If the atoms of H are smaller than the atoms of J, then compared to atoms of J, atoms of H are most likely to
  - (A) exist in a greater number of isotopes
  - (B) exist in a lesser number of isotopes
  - (C) exist in a greater number of oxidation states
  - (D) have a greater positive charge in their nuclei
  - (E) have a lesser positive charge in their nuclei

 $\dots$  Al(s) +  $\dots$  O<sub>2</sub>(g)  $\rightarrow$   $\dots$  Al<sub>2</sub>O<sub>3</sub>(s)

- 45. When the equation representing the reaction shown above is completed and balanced and all coefficients are reduced to lowest whole-number terms, the coefficient of  $O_2(g)$  is
  - (A) 1
  - (B) 2
  - (C) 3
  - (D) 4
  - (E) 6
- 46. Which of the following solids has a brilliant blue color?
  - (A)  $Ca(OH)_{2}$
  - (B) KCl
  - (C) NaBr
  - (D)  $Fe_2O_3$
  - (E)  $CuSO_4$
- 47. Twenty-five percent of element X exists as <sup>210</sup>X and 75 percent of it exists as <sup>214</sup>X. What is the atomic weight of element X in amu?
  - (A) 85
  - (B) 211
  - (C) 212
  - (D) 213
  - (E) 214



- 48. A 600-milliliter container holds 2 moles of  $O_2(g)$ , 3 moles of  $H_2(g)$ , and 1 mole of He(g). Total pressure within the container is 760 torr. What is the partial pressure of  $O_2$ ?
  - (A) 127 torr
  - (B) 253 torr
  - (C) 380 torr
  - (D) 507 torr
  - (E) 760 torr

 $\operatorname{Fe(OH)}_{3}(s) \sqsubseteq \operatorname{Fe}^{3+}(aq) + \operatorname{3OH}^{-}(aq)$ 

- 49. The ionic solid  $\text{Fe(OH)}_3$  is added to water and dissociates into its component ions, as shown above. The solubility product expression for the saturated solution is
  - (A)  $K_{sp} = [Fe^{3+}] [OH^{-}]$
  - (B)  $K_{sp} = [Fe^{3+}] [3OH^{-}]$
  - (C)  $K_{sp} = [Fe^{3+}] [3OH^{-}]^{3}$
  - (D)  $K_{sp} = [Fe^{3+}] [OH^{-}]^3$ (E)  $K_{sp} = \frac{[Fe^{3+}] [OH^{-}]^3}{[Fe(OH)_2]}$
- 50. Which of the following electron configurations represents an atom of magnesium in an excited state?
  - (A)  $1s^2 2s^2 2p^6$
  - (B)  $1s^22s^22p^63s^2$
  - (C)  $1s^22s^22p^53s^23p^2$
  - (D)  $1s^22s^22p^63s^13p^1$
  - (E)  $1s^22s^22p^63s^13p^2$

- 51. All of the following when added to water will produce an electrolytic solution EXCEPT
  - (A)  $N_{2}(g)$
  - (B) HCl(g)
  - (C) KOH(*s*)
  - (D) NaI(s)
  - (E)  $\operatorname{CaCl}_2(s)$

 $NH_3(aq) + H_2CO_3(aq) \rightleftharpoons NH_4^+(aq) + HCO_3^-(aq)$ 

- 52. In the reaction represented above,  $NH_4^+$  acts as
  - a(n)
  - (A) indicator
  - (B) hydrate
  - (C) acid
  - (D) base
  - (E) salt



- 53. Which species has the ground state electron configuration  $1s^22s^22p^63s^23p^6$ ?
  - (A) Sulfide ion, S<sup>2-</sup>
  - (B) Bromide ion, Br-
  - (C) Neon atom, Ne
  - (D) Chromium ion,  $Cr^{3+}$
  - (E) Potassium atom, K

- 54. Which of the following species is amphoteric?
  - (A)  $Na_3PO_4$
  - (B)  $HSO_4^-$
  - (C) KOH
  - (D) HNO<sub>3</sub>
  - (E)  $C_2 O_4^{2-}$
- 55. An ideal gas has a volume of 10 liters at 20° C and a pressure of 750 mmHg. Which of the following expressions is needed to determine the volume of the same amount of gas at STP?
  - (A)  $10 \times \frac{750}{760} \times \frac{0}{20}$  L

(B) 
$$10 \times \frac{750}{760} \times \frac{293}{273}$$
 L

- (C)  $10 \times \frac{760}{750} \times \frac{0}{20}$  L
- (D)  $10 \times \frac{760}{750} \times \frac{273}{293}$  L
- (E)  $10 \times \frac{750}{760} \times \frac{273}{293}$  L



**Questions 56–57** pertain to the phase diagram for substance Z below.



- 56. Substance Z is at 0.5 atm and 200 K. If the pressure on substance Z is steadily increased and its temperature is kept constant, what phase change will eventually occur?
  - (A) Condensation
  - (B) Freezing
  - (C) Melting
  - (D) Sublimation
  - (E) Vaporization
- 57. The normal boiling point of substance Z is closest to
  - (A) 100 K
  - (B) 200 K
  - (C) 300 K
  - (D) 400 K
  - (E) 500 K

- 58. The shape of a PCl<sub>3</sub> molecule is described as
  - (A) bent
  - (B) trigonal pyramidal
  - (C) linear
  - (D) trigonal planar
  - (E) tetrahedral
- 59. What volume of  $0.4 M \operatorname{Ba(OH)}_2(aq)$  is needed to exactly neutralize 100 milliliters of  $0.2 M \operatorname{HBr}(aq)$ ?
  - (A) 25 mL
  - (B) 50 mL
  - (C) 100 mL
  - (D) 200 mL
  - (E) 400 mL
- 60. Which of the following is true regarding the aqueous dissociation of HCN,  $K_a = 4.9 \times 10^{-10}$  at 25°C ?
  - I. At equilibrium,  $[H^+] = [CN^-]$
  - II. At equilibrium,  $[H^+] > [HCN]$
  - III. HCN(aq) is a strong acid.
  - (A) I only
  - (B) II only
  - $(C) \ \ I \ and \ II \ only$
  - (D) II and III only
  - (E) I, II, and III
- 61. Which of the following atoms has the largest second ionization energy?
  - (A) Silicon, Si
  - (B) Calcium, Ca
  - (C) Chlorine, Cl
  - (D) Iron, Fe
  - (E) Sodium, Na



**Question 62** refers to the overall reaction and halfreactions with standard reduction potentials below.

$$2Fe^{2+} + Cl_2 \rightarrow 2Fe^{3+} + 2Cl^{-}$$
  
Fe<sup>3+</sup> + e<sup>-</sup>  $\rightarrow$  Fe<sup>2+</sup>;  $E^{o}_{red} = 0.77$  volts  
Cl<sub>2</sub> + 2e<sup>-</sup>  $\rightarrow$  2Cl<sup>-</sup>;  $E^{o}_{red} = 1.36$  volts

- 62. The standard potential difference of an electrochemical cell using the overall reaction above is
  - (A) 0.18 volts
  - (B) 0.59 volts
  - (C) 1.05 volts
  - (D) 2.13 volts
  - (E) 2.90 volts
- 63. The reaction of zinc metal, Zn, and hydrochloric acid, HCl, produces which of the following?
  - I.  $H_2(g)$
  - II.  $\overline{Cl}_{2}(g)$
  - III.  $Zn^{2+}(aq)$
  - (A) II only
  - (B) III only
  - (C) I and II only
  - (D) I and III only
  - (E) I, II, and III

Questions 64–65 refer to the following reaction.

 $2H_2S(g) + 3O_2(g) \rightleftharpoons 2SO_2(g) + 2H_2O(g) + heat$ 

- 64. For the above reaction, the equilibrium concentration of  $SO_2(g)$  can be increased by
  - (A) adding neon gas
  - (B) increasing the temperature
  - (C) adding a catalyst
  - (D) increasing the concentration of  $H_2O(g)$
  - (E) increasing the concentration of  $O_2(g)$
- 65. Which of the following is increased by decreasing the volume of the reaction system?
  - I. Rate of reaction
  - II. Equilibrium concentration of reactants
  - III. Value of  $K_{ea}$
  - (A) I only
  - (B) III only
  - (C) I and II only
  - (D) II and III only
  - (E) I, II, and III

 $\operatorname{Fe}_{2}\operatorname{O}_{3}(s) + 3\operatorname{CO}(g) \rightarrow 2\operatorname{Fe}(s) + 3\operatorname{CO}_{2}(g)$ 

- 66. When 3 moles of  $Fe_2O_3$  are allowed to completely react with 56 grams of CO according to the above equation, approximately how many moles of iron, Fe, are produced?
  - (A) 0.7
  - (B) 1.3
  - (C) 2.0
  - (D) 2.7(E) 6.0



 $2\mathrm{Na}_{2}\mathrm{O}_{2}(s) + 2\mathrm{H}_{2}\mathrm{O}(l) \rightarrow 4\mathrm{NaOH}(aq) + \mathrm{O}_{2}(g)$ 

- 67. Sodium peroxide,  $Na_2O_2$ , and water react in the flask at 25°C according to the equation and in the diagram above. If water levels are equal inside and outside the beaker, then the gas pressure inside the beaker is equal to the
  - (A) pressure of oxygen gas collected
  - (B) vapor pressure of water at 25°C
  - (C) sum of pressure of oxygen gas collected and atmospheric pressure
  - (D) sum of vapor pressure of water at 25°C and atmospheric pressure
  - (E) sum of pressure of oxygen gas collected and vapor pressure of water at 25°C
- 68. Which of the following molecules has the strongest carbon-to-carbon bond?
  - (A)  $C_2H_2$
  - (B)  $C_2H_4$
  - $(C) C_2H_2$
  - (D)  $C_3H_8$
  - (E)  $C_4 H_{10}$

 $N_2O_4(g) \rightleftharpoons 2NO_2(g)$ 

The following concentration data were gathered for the above reaction at 5 minute intervals from the start of an experiment:

#### Time After Start

| of Experiment | $[N_2O_4]$ | $[NO_2]$ |  |  |  |
|---------------|------------|----------|--|--|--|
| 0 min (start) | 0.00 M     | 0.50 M   |  |  |  |
| 5 min         | 0.10 M     | 0.33 M   |  |  |  |
| 10 min        | 0.20 M     | 0.20 M   |  |  |  |
| 15 min        | 0.25 M     | 0.15 M   |  |  |  |
| 20 min        | 0.28 M     | 0.13 M   |  |  |  |
| 25 min        | 0.28 M     | 0.13 M   |  |  |  |
|               |            |          |  |  |  |

- 69. If the experiment was carried out in a closed system at constant temperature, then during which time interval (from the start of the experiment) did the reaction most likely achieve equilibrium?
  - (A) 0 min (start) to 5 min
  - (B)  $5 \min to 10 \min$
  - (C) 10 min to 15 min
  - (D) 15 min to 20 min
  - (E) 20 min to 25 min

# STOP

If you finish before time is called, you may check your work on this section only. Do not turn to any other section in the test.