

Practice Test 2

AP[®] Physics 1 Exam

SECTION I: Multiple-Choice Questions

DO NOT OPEN THIS BOOKLET UNTIL YOU ARE TOLD TO DO SO.

At a Glance

Total Time 90 minutes Number of Questions 50 Percent of Total Grade 50% Writing Instrument Pen required

Instructions

Section I of this examination contains 50 multiple-choice questions. Fill in only the ovals for numbers 1 through 50 on your answer sheet.

CALCULATORS MAY BE USED ON BOTH SECTIONS OF THE AP PHYSICS 1 EXAM.

Indicate all of your answers to the multiple-choice questions on the answer sheet. No credit will be given for anything written in this exam booklet, but you may use the booklet for notes or scratch work. Please note that there are two types of multiple-choice questions: single-select and multi-select questions. After you have decided which of the suggested answers is best, completely fill in the corresponding oval(s) on the answer sheet. For single-select, you must give only one answer; for multi-select you must give BOTH answers in order to earn credit. If you change an answer, be sure that the previous mark is erased completely. Here is a sample question and answer.

Sample Question



 $(A) \odot (D)$

Chicago is a (A) state

- (B) city
- (C) country
- (D) continent

Use your time effectively, working as quickly as you can without losing accuracy. Do not spend too much time on any one question. Go on to other questions and come back to the ones you have not answered if you have time. It is not expected that everyone will know the answers to all the multiple-choice questions.

About Guessing

Many candidates wonder whether or not to guess the answers to questions about which they are not certain. Multiple-choice scores are based on the number of questions answered correctly. Points are not deducted for incorrect answers, and no points are awarded for unanswered questions. Because points are not deducted for incorrect answers, you are encouraged to answer all multiple-choice questions. On any questions you do not know the answer to, you should eliminate as many choices as you can, and then select the best answer among the remaining choices.

Section I

ADVANCED PLACEMENT PHYSICS 1 TABLE OF INFORMATION

CONSTANTS AND CONVERSION FACTORS						
Proton mass, $m_p = 1.67 \times 10^{-27}$ kg	Electron charge magnitude, $e = 1.60 \times 10^{-19} \text{ C}$					
Neutron mass, $m_n = 1.67 \times 10^{-27}$ kg	Coulomb's law constant, $k = 1/4\pi\varepsilon_0 = 9.0 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$					
Electron mass, $m_e = 9.11 \times 10^{-31} \text{ kg}$	Universal gravitational constant, $G = 6.67 \times 10^{-11} \text{ m}^3/\text{kg} \cdot \text{s}^2$					
Speed of light, $c = 3.00 \times 10^8 \text{ m/s}$	Acceleration due to gravity at Earth's surface, $g = 9.8 \text{ m/s}^2$					

	meter,	m	kelvin,	Κ	watt,	W	degree Celsius,	°C
UNIT	kilogram,	kg	hertz,	Hz	coulomb,	С		
SYMBOLS	second,	S	newton,	Ν	volt,	V		
	ampere,	А	joule,	J	ohm,	Ω		

	PREFIXES						
Factor	Factor Prefix						
10 ¹²	tera	Т					
10 ⁹	giga	G					
10 ⁶	mega	М					
10 ³	kilo	k					
10 ⁻²	centi	с					
10 ⁻³	milli	m					
10 ⁻⁶	micro	μ					
10 ⁻⁹	nano	n					
10 ⁻¹²	pico	р					

VALUES OF TRIGONOMETRIC FUNCTIONS FOR COMMON ANGLES								
θ	0°	30°	37°	45°	53°	60°	90°	
sin $ heta$	0	1/2	3/5	$\sqrt{2}/2$	4/5	$\sqrt{3}/2$	1	
$\cos \theta$	1	$\sqrt{3}/2$	4/5	$\sqrt{2}/2$	3/5	1/2	0	
tan 0	0	$\sqrt{3}/3$	3/4	1	4/3	$\sqrt{3}$	~	

The following conventions are used in this exam.

- I. The frame of reference of any problem is assumed to be inertial unless otherwise stated.
- II. Assume air resistance is negligible unless otherwise stated.
- III. In all situations, positive work is defined as work done <u>on</u> a system.
- IV. The direction of current is conventional current: the direction in which positive charge would drift.
- V. Assume all batteries and meters are ideal unless otherwise stated.

ADVANCED PLACEMENT PHYSICS 1 EQUATIONS, EFFECTIVE 2015

MECI	HANICS	ELEC	TRICITY
MECH $v_x = v_{x0} + a_x t$ $x = x_0 + v_{x0}t + \frac{1}{2}a_x t^2$ $v_x^2 = v_{x0}^2 + 2a_x(x - x_0)$ $\vec{a} = \frac{\sum \vec{F}}{m} = \frac{\vec{F}_{net}}{m}$ $ \vec{F}_f \le \mu \vec{F}_n $ $a_c = \frac{v^2}{r}$ $\vec{p} = m\vec{v}$ $\Delta \vec{p} = \vec{F} \Delta t$	$a = \operatorname{acceleration} A$ $A = \operatorname{amplitude} d$ $d = \operatorname{distance} E$ $E = \operatorname{energy} f$ $f = \operatorname{frequency} F$ $F = \operatorname{force} I$ $I = \operatorname{rotational inertia} K$ $K = \operatorname{kinetic energy} k$ $k = \operatorname{spring constant} L$ $L = \operatorname{angular momentum} \ell$ $\ell = \operatorname{length} m$ $m = \max S$ $P = \operatorname{power} p$ $p = \operatorname{momentum} r$ $r = \operatorname{radius or separation} T$ $T = \operatorname{period} t$ $t = \operatorname{time} $	ELEC $\begin{aligned} \vec{F}_E &= k \left \frac{q_1 q_2}{r^2} \right \\ I &= \frac{\Delta q}{\Delta t} \\ R &= \frac{\rho \ell}{A} \\ I &= \frac{\Delta V}{R} \\ P &= I \Delta V \\ R_s &= \sum_i R_i \\ \frac{1}{R_p} &= \sum_i \frac{1}{R_i} \end{aligned}$	FRICTLY A = area F = force I = current $\ell = \text{length}$ P = power q = charge R = resistance r = separation t = time V = electric potential $\rho = \text{resistivity}$
$K = \frac{1}{2}mv^2$	U = potential energy V = volume	W	AVES
$\Delta E = W = F_{\parallel}d = Fd\cos\theta$	v = volume v = speed W = work done on a system x = position	$\lambda = \frac{v}{f} \qquad \qquad f = v = v$	frequency
$P = \frac{\Delta E}{\Delta t}$	y = height $\alpha =$ angular acceleration		D TRIGONOMETRY
$\theta = \theta_0 + \omega_0 t + \frac{1}{2}\alpha t^2$	μ = coefficient of friction θ = angle ρ = density	Rectangle $A = bh$	A = area C = circumference V = volume
$\omega = \omega_0 + \alpha t$	τ = torque	Triangle $A = \frac{1}{2} hh$	S = surface area b = base
$x = A\cos(2\pi ft)$ $\vec{\alpha} = \frac{\sum \vec{\tau}}{I} = \frac{\vec{\tau}_{net}}{I}$ $\tau = r_{\perp}F = rF\sin\theta$	ω = angular speed $\Delta U_g = mg \Delta y$ $T = \frac{2\pi}{\omega} = \frac{1}{f}$	$A = \frac{1}{2}bh$ Circle $A = \pi r^{2}$ $C = 2\pi r$	h = height $\ell = length$ w = width r = radius
$L = I\omega$ $\Delta L = \tau \Delta t$	ωf $T_s = 2\pi \sqrt{\frac{m}{k}}$	Rectangular solid $V = \ell w h$	Right triangle $c^2 = a^2 + b^2$
$K = \frac{1}{2}I\omega^2$ $\left \vec{F}_s\right = k\left \vec{x}\right $	$T_p = 2\pi \sqrt{\frac{\ell}{g}}$	Cylinder $V = \pi r^2 \ell$ $S = 2\pi r \ell + 2\pi r^2$	$\sin \theta = \frac{a}{c}$ $\cos \theta = \frac{b}{c}$
$U_s = \frac{1}{2}kx^2$ $\rho = \frac{m}{V}$	$\left \vec{F}_{g}\right = G \frac{m_{1}m_{2}}{r^{2}}$ $\vec{g} = \frac{\vec{F}_{g}}{m}$	Sphere $V = \frac{4}{3}\pi r^{3}$ $S = 4\pi r^{2}$	$\tan \theta = \frac{a}{b}$ c $\theta = 90^{\circ}$
v	$U_G = -\frac{Gm_1m_2}{r}$		b

AP PHYSICS 1


SECTION I

Note: To simplify calculations, you may use $g = 10 \text{ m/s}^2$ in all problems.


Directions: Each of the questions or incomplete statements is followed by four suggested answers or completions. Select the one that is best in each case and then fill in the corresponding circle on the answer sheet.

1. An object is thrown horizontally to the right off a high cliff with an initial speed of 7.5 m/s. Which arrow best represents the direction of the object's velocity after 2 seconds? (Assume air resistance is negligible.)






Questions 2-4 refer to the following figure:




- 2. Which of the following correctly ranks the change in kinetic energy for each segment from least to greatest?
 - (A) BC, EF, DE, CD, AB
 - (B) AB, CD, DE, EF, BC
 - (C) BC, EF, DE, CD, AB
 - (D) CD, DE, EF, BC, AB

- 3. During which segment is the magnitude of average acceleration greatest?
 - (A) AB
 - (B) BC
 - (C) CD
 - (D) DE
- 4. What is the total distance traveled by the object?
 - (A) 32 m
 - (B) 34 m
 - (C) 36 m
 - (D) 38 m
- 5. A ball is thrown in a projectile motion trajectory with an initial velocity v at an angle θ above the ground. If the acceleration due to gravity is -g, which of the following is the correct expression of the time it takes for the ball to reach its highest point, y, from the ground?
 - (A) $v^2 \sin t/g$
 - (B) $-v \cos \theta/g$
 - (C) $v \sin \theta/g$
 - (D) $v^2 \cos \theta/g$
- 6. A bubble in a glass of water releases from rest at the bottom of the glass and rises at acceleration *a* to the surface in *t* seconds. How much farther does the bubble travel in its last second than in its first second?
 - (A) at
 - (B) (t-1)a
 - (C) (t+1)a
 - (D) $\frac{1}{2}at$


- 7. A person standing on a horizontal floor is acted upon by two forces: the downward pull of gravity and the upward normal force of the floor. These two forces
 - (A) have equal magnitudes and form an action-reaction pair
 - (B) have equal magnitudes and do not form an actionreaction pair
 - (C) have unequal magnitudes and form an actionreaction pair
 - (D) have unequal magnitudes and do not form an actionreaction pair
- 8. Which of the following graphs best represents the force of friction on an object starting at rest that eventually starts sliding across a level surface due to a gradually increasing horizontal force?



- 9. Two objects have a mass 1 kg and carry a charge of magnitude 1 C each. Which statement correctly identifies the relationship between the magnitude of the force of gravity, F_g, and the magnitude of the electric force, F_E, between the objects?
 - (A) $F_g > F_E$
 - (B) $F_g < F_E$
 - (C) $F_g = F_E$
 - (D) Cannot be determined without knowing the sign of the charges.



- 10. Consider the above configuration of masses attached via a massless rope and pulley over a frictionless inclined plane. What is the acceleration of the masses?
 - (A) $(m_1 m_2) g/(m_1 + m_2)$
 - (B) $(m_1 m_2 \sin \theta) g/(m_1 + m_2)$
 - (C) $(m_1 m_2 \cos \theta) g/(m_1 + m_2)$
 - (D) g
- 11. A person is pulling a block of mass *m* with a force equal to its weight directed 30° above the horizontal plane across a rough surface, generating a friction *f* on the block. If the person is now pushing downward on the block with the same force 30° above the horizontal plane across the same rough surface, what is the friction on the block? (μ_k is the coefficient of kinetic friction across the surface.)
 - (A) f(B) 1.5f
 - (C) 2f
 - (D) 3f




- 12. In the figure above, two blocks of mass 3m and 2m are attached together. The plane is frictionless and the pulley is frictionless and massless. The inclined portion of the plane creates an angle θ with the horizontal floor. What is the acceleration of the block 2m if both blocks are released from rest (gravity = g)?
 - (A) 2mg


(B)
$$\left(\frac{2}{5}\right)g\sin\theta$$

(C) $\left(\frac{2}{3}\right)g\sin\theta$
(D) $\left(\frac{3}{5}\right)g\sin\theta$

- 13. If a roller coaster cart of mass *m* was not attached to the track, it would still remain in contact with a track throughout a loop of radius *r* as long as
 - (A) $v \leq \sqrt{(rg)}$
 - (B) $v \ge \sqrt{(rg)}$
 - (C) $v \leq \sqrt{(rg/m)}$
 - (D) $v \ge \sqrt{(rg/m)}$




- 14. The diagram above shows a top view of an object of mass *M* on a circular platform of mass 2*M* that is rotating counterclockwise. Assume the platform rotates without friction. Which of the following best describes an action by the object that will increase the angular speed of the entire system?
 - (A) The object moves toward the center of the platform, increasing the total angular momentum of the system.
 - (B) The object moves toward the center of the platform, decreasing the rotational inertia of the system.
 - (C) The object moves away from the center of the platform, increasing the total angular momentum of the system.
 - (D) The object moves away from the center of the platform, decreasing the rotational inertia of the system.



- 15. A moon has an elliptical orbit about the planet as shown above. At point *A*, the moon has speed v_A and is at a distance r_A from the planet. At point *B*, the moon has a speed of v_B . Which of the following correctly explains the method for determining the distance of the moon from the planet at point *B* in the given quantities?
 - (A) Conservation of angular momentum, because the gravitational force exerted by the moon on the planet is the same as that exerted by the planet on the moon
 - (B) Conservation of angular momentum, because the gravitational force exerted on the moon is always directed toward the planet
 - (C) Conservation of energy, because the gravitational force exerted on the moon is always directed toward the planet
 - (D) Conservation of energy, because the gravitational force exerted by the moon on the planet is the same as that exerted by the planet on the moon

- 16. A sphere starts from rest atop a hill with a constant angle of inclination and is allowed to roll without slipping down the hill. What force provides the torque that causes the sphere to rotate?
 - (A) Static friction
 - (B) Kinetic friction
 - (C) The normal force of the hill on the sphere
 - (D) Gravity
- 17. Which of the following correctly describes the motion of a real object in free fall? Assume that the object experiences drag force proportional to speed and that it strikes the ground before reaching terminal sped.
 - (A) It will fall with increasing speed and increasing acceleration.
 - (B) It will fall with increasing speed and decreasing acceleration.
 - (C) It will fall with decreasing speed and increasing acceleration.
 - (D) It will fall with decreasing speed and decreasing acceleration.
- 18. Which of the following concerning uniform circular motion is true?
 - (A) The centrifugal force is the action-reaction pair of the centripetal force.
 - (B) The centripetal acceleration and velocity point in the same direction.
 - (C) The velocity of the object in motion changes whereas the acceleration of the object is constant.
 - (D) A satellite undergoing uniform circular motion is falling toward the center in a circular path.
- 19. A girl of mass *m* and a boy of mass 2*m* are sitting on opposite sides of a see-saw with its fulcrum in the center. Right now, the boy and girl are equally far from the fulcrum, and it tilts in favor of the boy. Which of the following would NOT be a possible method to balance the see-saw?
 - (A) Move the boy to half his original distance from the fulcrum.
 - (B) Move the girl to double her original distance from the fulcrum.
 - (C) Allow a second girl of mass *m* to join the first.
 - (D) Move the fulcrum to half its original distance from the boy.

- 20. Given that the Earth's mass is *m*, its tangential speed as it revolves around the Sun is *v*, and the distance from the Sun to the Earth is *r*, which of the following correctly describes the work done by the centripetal force, W_c , in one year's time?
 - (A) $W_c > 2r(mv^2/r)$
 - (B) $W_c = 2r(mv^2/r)$
 - (C) $W_c < 2r(mv^2/r)$
 - (D) Cannot be determined
- 21. A wooden block experiences a frictional force, *f*, as it slides across a table. If a block of the same material with half the height and twice the length were to slide across the table, what would be the frictional force it experienced?
 - (A) (1/2)f
 - (B) *f*
 - (C) 2f
 - (D) 4f
- 22. Two objects, a sphere and a block of the same mass, are released from rest at the top of an inclined plane. The sphere rolls down the inclined plane without slipping. The block slides down the plane without friction. Which object reaches the bottom of the ramp first?
 - (A) The sphere, because it gains rotational kinetic energy, but the block does not
 - (B) The sphere, because it gains mechanical energy due to the torque exerted on it, but the block does not
 - (C) The block, because it does not lose mechanical energy due to friction, but the sphere does
 - (D) The block, because it does not gain rotational kinetic energy, but the sphere does




- 23. In the diagram above, a mass *m* starting at point *A* is projected with the same initial horizontal velocity v_0 along each of the three tracks shown here (with negligible friction) sufficient in each case to allow the mass to reach the end of the track at point *B*. (Path 1 is directed up, path 2 is directed horizontal, and path 3 is directed down.) The masses remain in contact with the tracks throughout their motions. The displacement *A* to *B* is the same in each case, and the total path length of path 1 and 3 are equal. If t_1 , t_2 , and t_3 are the total travel times between *A* and *B* for paths 1, 2, and 3, respectively, what is the relation among these times?
 - $(A) \quad t_3 < t_2 < t_1$


(B)
$$t_2 < t_3 < t_1$$

- (C) $t_2 < t_1 = t_3$
- (D) $t_2 = t_3 < t_1$
- 24. An object of mass m_1 experiences a linear, elastic collision with a stationary object of unknown mass. In addition to m_1 , what is the minimum necessary information that would allow you to determine the mass of the second object?
 - (A) The final speed of object 1
 - (B) The initial speed of object 1
 - (C) The final speed of object 2
 - (D) Any 2 of the above values
- 25. A block is dragged along a table and experiences a frictional force, f, that opposes its movement. The force exerted on the block by the table is
 - (A) zero
 - (B) parallel to the table
 - (C) perpendicular to the table
 - (D) neither parallel nor perpendicular to the table


26. A box of mass *m* is sitting on an incline of 45° and it requires an applied force *F* up the incline to get the box to begin to move. What is the maximum coefficient of static friction?



- 27. The graph above shows the velocities of two objects undergoing a head-on collision. Given that Object 1 has 4 times the mass of Object 2, which type of collision is it?
 - (A) Perfectly elastic
 - (B) Perfectly inelastic
 - (C) Inelastic
 - (D) Cannot be determined



28. The picture above depicts the collision of two balls of equal mass. Which arrow best indicates the direction of the impulse on ball A from ball B during the collision?



- 29. Which of the following best describes the relationship between the magnitude of the tension force, F_{T} , in the string of a pendulum and the radial component of gravity that pulls antiparallel to the tension, $F_{\rm g, radial}$? Assume that the pendulum is only displaced by a small amount.
 - (A) $F_{\rm T} > F_{\rm g, radial}$
 - (B) $F_{\rm T} \ge F_{\rm g, radial}$
 - (C) $F_{\rm T} = F_{\rm g, radial}$ (D) $F_{\rm T} \le F_{\rm g, radial}$


- 30. Pretend someone actually managed to dig a hole straight through the center of the Earth all the way to the other side. If an object were dropped down that hole, which of the following would best describe its motion? Assume ideal conditions and that the object cannot be destroyed.
 - (A) It would fall to the center of the Earth and stop there.
 - (B) It would fall through the hole to the other side, continue past the opposite side's opening, and fly into space.
 - (C) It would oscillate back and forth from one opening to the other indefinitely.
 - (D) It would fall to the other side and stop there.
- 31. A sound wave with frequency f travels through air at speed v. With what speed will a sound wave with frequency 4f travel through the air?
 - (A) v/4
 - (B) v
 - (C) 2v
 - (D) 4v
- 32. An object has a kinetic energy of K. If the speed of the object were doubled what would be the resulting kinetic energy?
 - (A) $\frac{1}{2}K$
 - (B) *K*
 - (C) 2K
 - (D) 4K
- 33. A toy car and a toy truck collide. If the toy truck's mass is double the toy car's mass, then, compared to the acceleration of the truck, the acceleration of the car during the collision will be
 - (A) double the magnitude and in the same direction
 - (B) double the magnitude and in the opposite direction
 - (C) half the magnitude and in the same direction
 - (D) half the magnitude and in the opposite direction

Section I

- 34. The Gravitron is a carnival ride that looks like a large cylinder. People stand inside the cylinder against the wall as it begins to spin. Eventually, it is rotating fast enough that the floor can be removed without anyone falling. Given then the coefficient of friction between a person's clothing and the wall is μ , the tangential speed is v, and the radius of the ride is r, what is greatest mass that a person can be to safely go on this ride?
 - (A) $\mu v^2/(rg)$
 - (B) $r^2 v^2 / (\mu g)$
 - (C) $rg/(\mu v^2)$
 - (D) None of the above.
- 35. In a spring-block oscillator, the maximum speed of the block is
 - (A) proportional to amplitude
 - (B) proportional to the square of amplitude
 - (C) proportional to the square root of amplitude
 - (D) inversely proportional to the square root of amplitude
- 36. A student is experimenting with a simple spring-block oscillator of spring constant *k* and amplitude *A*. The block attached to the spring has a mass of *M*. If the student places a small block of mass *m* on top of the original block, which of the following is true?
 - (A) The small block is most likely to slide off when the original block is at maximum displacement from the equilibrium position, but will not slide off as long as the coefficient of static friction between the blocks is greater than kA/[(M+m)g].
 - (B) The small block is most likely to slide off when the original block is at the equilibrium position, but will not slide off as long as the coefficient of static friction between the blocks is greater than kA/[(M+m)g].
 - (C) The small block is most likely to slide off when the original block is at maximum displacement from the equilibrium position, but will not slide off as long as the coefficient of static friction between the blocks is greater than (M+m)g/(kA).
 - (D) The small block is most likely to slide off when the original block is at the equilibrium position, but will not slide off as long as the coefficient of static friction between the blocks is greater than (M+m)g/(kA).

- 37. A pair of protective headphones reduces the sound of a jet engine from 140 dB to 110 dB. What percent of the noise is being blocked?
 - (A) 99.9%
 - (B) 78.6%
 - (C) 21.4%
 - (D) 0.1%
- 38. You are standing on a railroad track as a train approaches at a constant velocity. Suddenly the engineer sees you, applies the brakes, and sounds the whistle. Which of the following describes the sound of the whistle as you hear it starting from that moment?
 - (A) Loudness increasing, pitch increasing
 - (B) Loudness increasing, pitch constant
 - (C) Loudness decreasing, pitch increasing
 - (D) Loudness increasing, pitch decreasing

Questions 39-41 refer to the following figure:



39. Determine the total power dissipated through the circuit shown above in terms of V, R_1 , R_2 , and R_3 .

(A)
$$\frac{V^2}{R_1 + R_2 + R_3}$$

(B)
$$\frac{R_1 + R_2 + R_3}{V^2}$$

(C)
$$\frac{R_1(R_2 + R_3)}{V^2(R_1 + R_2 + R_3)}$$

(D)
$$\frac{V^2(R_1 + R_2 + R_3)}{R_1(R_2 + R_3)}$$


- 40. If V = 100 V, $R_1 = 50 \Omega$, $R_2 = 80 \Omega$ and $R_3 = 120 \Omega$, determine the voltage across R_3 .
 - (A) 100 V
 - (B) 60 V
 - (C) 40 V
 - (D) 20 V
- 41. If R_1 were to burn out, the current coming out from the battery would
 - (A) increase
 - (B) decrease
 - (C) stay the same
 - (D) there is no current, because the circuit is now incomplete
- 42. Wire #1 has three times the length and twice the diameter of Wire #2. If both wires are made of the same material and if R_1 is the resistance of Wire #1 and R_2 is the resistance of Wire #2, then which of the following is true?

(A)
$$R_2 = \left(\frac{2}{3}\right)R_1$$


(B) $R_2 = \left(\frac{4}{3}\right)R_1$

(C) $R_2 = 6R_1$

(D)
$$R_2 = 12R_1$$



- 43. A Wheatstone bridge (diagram above) is a configuration of resistors and a sensitive current meter, called a *gal-vanometer*, that is used to determine the resistance of an unknown resistor. In the Wheatstone bridge shown here, find the value of R_x such that the current through galvanometer *G* is zero.
 - (A) 25 Ω
 - (B) 15 Ω
 - (C) 10 Ω
 - (D) 2.5 Ω
- 44. If T, C, M, and L represent the dimensions of time, charge, mass, and length, respectively, which of the following represents the correct dimensions for the permittivity of free space (ε_0) ?
 - (A) $T^2C^2/(ML^3)$
 - (B) $ML^{3}/(T^{2}C^{2})$
 - (C) $C^2M/(T^2L^2)$
 - (D) $T^{2}L^{2}/(C^{2}M)$



45. In the figure above, four charges are arranged. If the magnitudes of all the charges *q* are all the same and the distance *r* between them is as shown above, what is the magnitude of the net force on the bottom right charge in terms of *q*, *r*, and *k* (where $k = \frac{1}{4\pi\varepsilon_0}$)?

(A)
$$k\left(\frac{q^2}{2r^2}\right)(1+\sqrt{2})$$

(B) $k\left(\frac{q^2}{r^2}\right)(1+\sqrt{2})$
(C) $k\left(\frac{q^2}{2r^2}\right)$
(D) $k\left(\frac{q^2}{r^2}\right)$

Directions: For each of the questions 46-50, <u>two</u> of the suggested answers will be correct. Select the two answers that are best in each case, and then fill in both of the corresponding circles on the answer sheet.

- 46. An object traveling at x m/s can stop at a distance d m with a maximum negative acceleration. If the car is traveling at 2x m/s, which of the following statements are true? Select two answers.
 - (A) The stopping time is doubled.
 - (B) The stopping time is quadrupled.
 - (C) The stopping distance is doubled.
 - (D) The stopping distance is quadrupled.

- 47. A 2 kg mass is attached to a massless, 0.5 m string and is used as a simple pendulum by extending it to an angle $\theta = 5^{\circ}$ and allowing it to oscillate. Which of the following changes will change the period of the pendulum? Select two answers.
 - (A) Replacing the mass with a 1 kg mass
 - (B) Changing the initial extension of the pendulum to a 10° angle
 - (C) Replacing the string with a 0.25 m string
 - (D) Moving the pendulum to the surface of the Moon
- 48. *N* resistors (N > 2) are connected in parallel with a battery of voltage V_o . If one of the resistors is removed from the circuit, which of the following quantities will decrease? Select two answers.
 - (A) The voltage across any of the remaining resistors
 - (B) The current output by the battery
 - (C) The total power dissipated in the circuit
 - (D) The voltage supplied by the battery
- 49. Which of the following statements will increase the resistance of a closed circuit system? Select two answers.
 - (A) Replacing the wire with one that has a smaller cross-section area
 - (B) Adding a voltmeter to the wire
 - (C) Halving the wire length in the close circuit
 - (D) Adding another resistor in parallel with the current resistor in a closed circuit
- 50. A sound wave travels through a metal rod with wavelength λ and frequency *f*. Which of the following is true? Select two answers.
 - (A) When this sound wave passes into air, the frequency will change.
 - (B) When this sound wave passes into air, the wavelength will change.
 - (C) While in the metal rod, the λ and frequency f have a direct relationship.
 - (D) While in the metal rod, the λ and frequency *f* have an inverse relationship.

END OF SECTION I

DO NOT CONTINUE UNTIL INSTRUCTED TO DO SO.

AP PHYSICS 1 SECTION II Free-Response Questions Time—90 minutes Percent of total grade—50


General Instructions

Use a separate piece of paper to answer these questions. Show your work. Be sure to write CLEARLY and LEGIBLY. If you make an error, you may save time by crossing it out rather than trying to erase it.


AP PHYSICS 1

SECTION II

Directions: Questions 1, 2, and 3 are short free-response questions that require about 13 minutes to answer and are worth 8 points. Questions 4 and 5 are long free-response questions that require about 25 minutes each to answer and are worth 13 points each. Show your work for each part in the space provided after that part.




- 1. An experiment is conducted in which Block A with a mass of m_A is slid to the right across a frictionless table. Block A collides with Block B, which is initially at rest, of an unknown mass and sticks to it.
 - (a) Describe an experimental procedure that determines the velocities of the blocks before and after a collision. Include all the additional equipment you need. You may include a labeled diagram of your setup to help in your description. Indicate what measurements you would take and how you would take them. Include enough detail so that the experiment could be repeated with the procedure you provide.
 - (b) If Block A has a mass of 0.5 kg and starts off with a speed of 1.5 m/s and the experiment is repeated, the velocity of the blocks after the collide are recorded to be 0.25 m/s. What is the mass of Block B?
 - (c) How much kinetic energy was lost in this collision from part (b)?




- 2. A conical pendulum is hanging from a string that is 2.2 meters long. It makes a horizontal circle. The mass of the ball at the end of the string is 0.5 kg.
 - (a) Below, make a free-body diagram for the ball at the point shown in the above illustration. Label each force with an appropriate letter.


- (b) Write out Newton's Second Law in both the *x* and *y*-direction in terms used in your free-body diagram.
- (c) Calculate the centripetal acceleration from your free-body diagram.
- (d) What is the radius of the circle that the ball is traveling in?
- (e) What is the speed of the ball?




- 3. Three particles are fixed in place in a horizontal plane, as shown in the figure above. Particle 1 has a charge that has a magnitude of 4.0×10^{-6} C and the sign of the charge is unknown. Particle 2 has a charge that has a magnitude of 1.7×10^{-6} C and the sign of the charge is unknown. Particle 3 has a charge of $+1.0 \times 10^{-6}$ C. The distance between q_1 and q_2 is 5.0 m, the distance between q_2 and q_3 is 3.0 m, and the distance between q_1 and q_3 is 4.0 m. The electrostatic force **F** on particle 3 due to the other two charges is shown in the negative *x*-direction.
 - (a) Determine the signs of the charges of q_1 and q_2 .
 - (b) On the diagram below, draw and label the force F_1 of the force exerted by Particle 1 on Particle 3 and the force F_2 of the force exerted by Particle 2 on Particle 3.



- (c) Calculate the magnitude of the electrostatic force on Particle 3.
- (d) Draw and label clearly where another positively charged particle could be placed so the net electrostatic force on Particle 3 is zero.






- 4. An experiment is designed to calculate the spring constant k of a vertical spring for a jumping toy. The toy is compressed a distance of x from its natural length of L_0 , as shown on the left in the diagram, and then released. When the toy is released, the top of the toy reaches a height of h in comparison to its previous height and the spring reaches its maximum extension. The experiment is repeated multiple times and replaced with different masses m attached to the spring. The spring itself has negligible mass.
 - (a) Derive an expression for the height h in terms of m, x, k, and any other constants provided in the formula sheet.
 - (b) To standardize the experiment, the compressed distance x is set to 0.020 m. The following table shows the data for different values of m.

<i>m</i> (kg)	<i>h</i> (m)
0.020	0.49
0.030	0.34
0.040	0.28
0.050	0.19
0.060	0.18


(i) What quantities should be graphed so that the slope of a best-fit straight line through the data points can help us calculate the spring constant k?

(ii) Fill in the blank column in the table above with calculated values. Also include a header with units.

(c) On the axes below, plot the data and draw the best-fit straight line. Label the axes and indicate scale.



- (d) Using your best-fit line, calculate the numerical value of the spring constant.
- (e) Describe an experimental procedure that determines the height h in the experiment, given that the toy is only momentarily at that maximum height. You may include a labeled diagram of your setup to help in your description.



- 5. In the above diagram, two small objects, each with a charge of -4.0 nC, are held together by a 0.020 m length of insulating string. The objects are initially at rest on a horizontal, non-conducting, frictionless surface. The effects of gravity on each other can be considered negligible.
 - (a) Calculate the tension in the string.
 - (b) Illustrate the electric field by drawing electric field lines for the two objects on the following diagram.



The masses of the objects are $m_1 = 0.030$ kg and $m_2 = 0.060$ kg. The string is now cut.

- (c) Calculate the magnitude of the initial acceleration of each object.
- (d) On the axes below, sketch a graph of the acceleration a of the object of mass m_2 versus the distance d between the objects after the string has been cut.



(e) In a brief paragraph, describe the speed of the objects as time increases, assuming that the objects remain on the horizontal, non-conducting frictionless surface.

STOP

Princeton Review [®]		Completely darken bubble	s with a No. 2 pencil. If	you make a	mistake, be sure to er	ase mark	complete	ely. Eras	e all stray ma	arks.	
1.						5. Y	OUR NA	ME			
YOUR NAME:		First		MI		First	4 letters	of last 1	name	FIRST	MID
SIGNATURE:			DATE:	/	/		A	A	A	A	A
						B	B	B	B	B	B
HOME ADDRESS:		Number and Street				\odot	Θ	\odot	\bigcirc	\odot	C
						\bigcirc	\bigcirc	\bigcirc	\bigcirc		
City	1	State		Zip Code	!	Ē		Ē	Ē	Ē	E
PHONE NO .						Ē	Ē	Ē	Ē	Ē	Ē
PHONE NO.:					_) G) (G	G	G	G	G
IMPORTANT: Please fill in these boxes exc	actly as shown on the back co	ver of your test book.				B B	9 8	Ð	Œ	Œ	(H
▼	· · · · · · · · · · · · · · · · · · ·	•				\square	Ē	\bigcirc	\square		
2. TEST FORM	3. TEST	CODE	4. REGISTRATION	NUMBER		Θ	9 (9	Ð	\bigcirc	
									_	_	
					\bigcirc	\mathbb{R}	$\Theta \Theta$	R	ß	ĸ	K
6. DATE OF BIRTH									_		
Month Day	v										
◯ JAN											
					33		Θ	\bigcirc	\odot		
						P	\mathbb{P}	P	P	P	P
		E O 5 5			5 5	0	0	0			Q
						R	R	®	R	R	R
		\mathbb{H}				S	9	S	S	S	S
					88	\square	Θ	\bigcirc	Ð		T
		99	9999	9	99	Θ	9	\bigcirc	\ominus		
						\bigotimes	\bigcirc	\bigcirc	\bigcirc	\bigtriangledown	
						\otimes	\otimes	W	(\mathbb{W})	\otimes	W
			The			\otimes	\bigotimes	\bigotimes	\bigotimes	\otimes	X
_	9 9 <u> </u>	EMALE	Princeto	n I		\heartsuit	$\langle \Sigma \rangle$	Y	\bigcirc	Y	Y
◯ DEC			Review			Z	\bigcirc	\bigcirc	\square	\square	Z
		I	INEVIEW								
1. A B C D E	_	BCDE							BC		
	-	$\mathbb{B} \mathbb{C} \mathbb{D} \mathbb{E}$									
3. A B C D E 4. A B C D E	_	BCDE							B C B C		
	-	BCDE							BC		
) 19. A (\mathbb{B} \mathbb{C} \mathbb{D} \mathbb{E}	32. A	$) \oplus ($	$\bigcirc \bigcirc \bigcirc \bigcirc$				BC		
7. A B C D E) 20. A ($\mathbb{B} \mathbb{C} \mathbb{D} \mathbb{E}$	_				46. (A	BC	DE	>
	<u> </u>	$\mathbb{B} \mathbb{C} \mathbb{D} \mathbb{E}$							BC		
	_	$\mathbb{B} \mathbb{C} \mathbb{D} \mathbb{E}$									
10. A B C D E 11. A B C D E	_	BCDE							B C B C		
	_	BCDE					JU. (
	<u> </u>	BCDE									