

Chapter 10 Drill

The answers and explanations can be found in Chapter 17.

Section I: Multiple Choice

- An object has an altitude of 2 times the Earth's radius, r_E , and experiences some force of gravity, $F_{g,0}$. If the object's altitude is doubled, then the new force of gravity will be
 - $4F_{g,0}$
 - $\left(\frac{25}{9}\right)F_{g,0}$
 - $2F_{g,0}$
 - $\left(\frac{1}{2}\right)F_{g,0}$
 - $\left(\frac{9}{25}\right)F_{g,0}$
- At the surface of Earth, an object of mass m has weight w . If this object is transported to a height above the surface that's twice the radius of Earth, then, at the new location,
 - its mass is $m/2$ and its weight is $w/2$
 - its mass is m and its weight is $w/2$
 - its mass is $m/2$ and its weight is $w/4$
 - its mass is m and its weight is $w/4$
 - its mass is m and its weight is $w/9$
- A moon of mass m orbits a planet of mass $100m$. Let the strength of the gravitational force exerted by the planet on the moon be denoted by F_1 , and let the strength of the gravitational force exerted by the moon on the planet be F_2 . Which of the following is true?
 - $F_1 = 100F_2$
 - $F_1 = 10F_2$
 - $F_1 = F_2$
 - $F_2 = 10F_1$
 - $F_2 = 100F_1$
- Humans cannot survive for long periods under gravity more than 4 times what we experience on Earth. If a planet were discovered with the same mass as Earth, what is the smallest radius it could have (in terms of Earth's radius, r_E) without being dangerous to humans?
 - $\frac{1}{8}r_E$
 - $\frac{1}{4}r_E$
 - $\frac{1}{2}r_E$
 - $2r_E$
 - $4r_E$
- Humans cannot survive for long periods under gravity more than 4 times what we experience on Earth. If a planet were discovered with the same mass as Earth, what is the smallest volume it could have (in terms of Earth's volume, V_E), without being dangerous to humans? Assume that the planet is a sphere.
 - $\frac{1}{8}V_E$
 - $\frac{1}{4}V_E$
 - $\frac{1}{2}V_E$
 - $2V_E$
 - $4V_E$
- A moon of Jupiter has a nearly circular orbit of radius R and an orbit period of T . Which of the following expressions gives the mass of Jupiter?
 - $2\pi R/T$
 - $4\pi^2 R/T^2$
 - $2\pi R^3/(GT^2)$
 - $4\pi R^2/(GT^2)$
 - $4\pi^2 R^3/(GT^2)$

7. Two large bodies, Body A of mass m and Body B of mass $4m$, are separated by a distance R . At what distance from Body A, along the line joining the bodies, would the gravitational force on an object be equal to zero? (Ignore the presence of any other bodies.)


- (A) $R/16$
 (B) $R/8$
 (C) $R/5$
 (D) $R/4$
 (E) $R/3$

8. The mean distance from Saturn to the Sun is 9 times greater than the mean distance from Earth to the Sun. How long is a Saturn year?

- (A) 18 Earth years
 (B) 27 Earth years
 (C) 81 Earth years
 (D) 243 Earth years
 (E) 729 Earth years

9. The Moon has mass M and radius R . A small object is dropped from a distance of $3R$ from the Moon's center. The object's impact speed when it strikes the surface of the Moon is equal to $\sqrt{kGM/R}$ for $k =$

- (A) $\frac{1}{3}$
 (B) $\frac{2}{3}$
 (C) $\frac{3}{4}$
 (D) $\frac{4}{3}$
 (E) $\frac{3}{2}$



10. Two satellites, A and B , orbit a planet in circular orbits having radii R_A and R_B , respectively, as shown above. If $R_B = 3R_A$, the velocities v_A and v_B of the two satellites are related by which of the following?

- (A) $v_B = v_A$
 (B) $v_B = 3v_A$
 (C) $v_B = 9v_A$
 (D) $v_B = v_A\sqrt{3}$
 (E) $v_B = \frac{v_A}{\sqrt{3}}$

Section II: Free Response


1. Consider two uniform spherical bodies in deep space. Sphere 1 has mass m_1 and Sphere 2 has mass m_2 . Starting from rest from a distance R apart, they are gravitationally attracted to each other.

- Compute the acceleration of Sphere 1 when the spheres are a distance $R/2$ apart.
- Compute the acceleration of Sphere 2 when the spheres are a distance $R/2$ apart.
- Compute the speed of Sphere 1 when the spheres are a distance $R/2$ apart.
- Compute the speed of Sphere 2 when the spheres are a distance $R/2$ apart.

Now assume that these spheres orbit their center of mass with the same orbit period, T .

- Determine the radii of their orbits. Write your answer in terms of m_1 , m_2 , T , and fundamental constants.

2. A satellite of mass m is in the elliptical orbit shown below around Earth (radius r_E , mass M). Assume that $m \ll M$.



- Determine v_1 , the speed of the satellite at perigee (the point of the orbit closest to Earth). Write your answer in terms of r_1 , r_2 , M , and G .
- Determine v_2 , the speed of the satellite at apogee (the point of the orbit farthest from Earth). Write your answer in terms of r_1 , r_2 , M , and G .
- Express the ratio v_1/v_2 in simplest terms.
- What is the satellite's angular momentum (with respect to Earth's center) when it's at apogee?
- Determine the speed of the satellite when it's at the point marked X in the figure.
- Determine the period of the satellite's orbit. Write your answer in terms of r_1 , r_2 , M , and fundamental constants.